



1ST AND 2ND ORDERS

The action of magnesium on dibromoethene and dibromo-*m*-xylene. V. S. ZALKIND, S. KIRILLOVA AND V. NIKIFOROV. *J. Gen. Chem. (U. S. S. R.)* 1, 103-8 (1931).—To a mixt. of 2.4 g. of Mg filings (previously washed with alc. and Et₂O and dried at 110°), a little dry Et₂O and a crystal of I was slowly added. 12.5 g. of 1,3,4-C₆H₄Br₂ (I) in 2 vols. of dry Et₂O and refluxed 35 hrs. in the water bath. On decompr. with ice water, the reaction mass was treated with 50 cc. of 0.8% H₂SO₄ to dissolve unchanged Mg, and the Br was measured, showing that 82.7% of Mg (and of Br) entered into reaction. The reaction mixt. was evap. with Et₂O and dried, giving 9 g. of resinous matter, 1.5 g. of PhMe, 3.5 g. of C₆H₄Mel₂ and 1.5 g. of high-boiling products. The tests showed that both Br atoms in I entered into the reaction with Mg, though incompletely and very slowly as compared with C₆H₄Br₂. Under similar conditions C₆H₄Mel₂ does not react with Mg and is completely recovered unchanged. When a mixt. of 2.4 g. of Mg filings, 13.2 g. of 1,3,4,6-C₆H₃Me₂Br₂, 3 vols. of dry Et₂O and a crystal of I are refluxed 60 hrs. and treated as described above, the reaction results in the consumption of 49.6% of Br with only 1 Br atom entering the reaction with Mg. Thus with increased no. of Me groups in the ring the reactivity of Br toward Mg is decreased.

Digitized by srujanika@gmail.com


436.34 METALLURGICAL LITERATURE CLASSIFICATION






"APPROVED FOR RELEASE: 09/19/2001


CIA-RDP86-00513R001963710009-5



APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"





Addition of hydrogen to acetylene derivatives. III.
Relation between the direction and progress of catalytic
hydrogenation and the nature of catalyst. Yu. S. Zelinskii.

Relation between the direction and progress of catalytic hydrogenation and the nature of catalyst. Yu. S. Zel'dovich, M. N. Vishnyakov and L. N. Morev, *J. Gen. Chem.* (U. S. S. R.) **3**, 91-113 (1933); *cit. C. A.* **25**, 3710. While the hydrogenation of acetylene γ -glycates (**I**) in the presence of Pd^0 black progresses uniformly with the formation of acid. glycate, the reaction with colloidal Pd proceeds in 2 stages: I rapidly adds 3 H atoms and then very slowly 2 more with the formation of acid. glycate. Such a sharp break in the process of hydrogenation is displayed only by γ -glycates. Acetylene also combines with 4 H atoms without retardation in the reaction, while acetylene hydrocyanoborane (**II**), such as $Pb(C_2H_5CH_3)_2$, adds the 2nd pair of H atoms even faster than the 1st (*C. A.* **8**, 1419; **10**, 1365; **17**, 1483). The hydrogenation of diphenylbutyryl acetate (**III**) also proceeds in 2 stages, with retardation of the reaction after addn. of 4 H atoms followed by highly increasing acceleration of the reduction with the addn. of 4 more H atoms and the formation of diphenylsuccinate. These facts indicate that the course of hydrogenation is determined not only by the speed, but that the reaction can proceed selectively, i.e., until the mols. of **II** ($Pb(C_2H_5CH_3)_2$) or those of an unsatd. glycol ester are present in the reaction system they will react with H₂ prior to other mols. (styrene or acid. glycol ester) available in the mixt., though separately the latter could combine with H₂ with a greater speed. Such a selective action of Pd was also demonstrated by Bourgat (*C. A.* **22**, 3040; **27**, 3372), who concluded that the selective action of


Although Pd suspended in starch always leads to hydrogenation of all available mols. of acetylene derivs. first to ethylene derivs. and thereafter to the mols. complexed. This is, action of Pd suspended in starch is not characteristic of other catalysts, such as Pt, or of Pt suspended in other protective colloids (Z., and Vilenkin, C. A. 10, 1461). Thus if the selective action of a catalyst is more definitely pronounced in some cases, it was of interest to inquire whether all acetylene derivs. act similarly, regardless of their structure, and to what extent this selective action is related to the chem. nature of the catalyst and is conditioned by the phys. form of the catalyst or the kind of protective colloid used for its suspension. In catalytic hydrogenation of acetylene derivs. there are formed *cis*- and *trans*-isomers of ethylene compds. While commonly the chief products are *trans*-isomeric isomers, the tetrasubhydrobenzene (IV) and *tetra*-phenylhydrobenzene produced also *summaric* isomers in various proportions, depending on the conditions of the reaction (Z., J. Russ. phys.-chem. Soc. 48, 1810 (1916); Z., and Teterin, C. A. 26, 78; Z., C. A. 28, 2450). Bourguel (C. A. 19, 2581; 24, 2451) ascribed the formation of the 2nd isomer of ethylene derived to some stereorearranging agents, such as protective colloids (protectant acids). IV, Me₂C(=O)₂ and PhC(=O)CH₃ were hydrogenated in the presence

ASA-SEA METALLURGICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: 09/19/2001


CIA-RDP86-00513R001963710009-5"

of Pd and Pt in the form of colloidal suspensions in protective colloids, powd. blacks and with various carriers. As protective colloids were used starch, gum arabic, gum tragacanth, agar, Na pyrolytinate and casein for Pd, and starch and gum arabic for Pt. The tabulated results of the exptl. work led to the following conclusions. The general course of hydrogenation of acetylene derivs. is conditioned by the chem. nature of the catalyst and the structure of the hydrogenated compd., and while different protective colloids and catalyst carriers may be unlike in their activity, they exert but little influence on the general progress of hydrogenation. The results substantiate the chem. character of the mechanism of catalysis. The catalyst displays selective affinity for hydrogenated unsat., showing in general a preference for unsat. with a triple bond. The character of catalytic distribution between acetylene and ethylene unsat. is conditioned by the nature of the catalyst and the structure of the compd. Thus Pd acts with more definite selectiveness than Pt, while HgCl₂CH₃ and styrene differ in their reactivity to a catalyst considerably more than acetylene does from the corresponding styrene alcs. All catalysts produced some α -tetramethylbutyrenediol by hydrogenation of IV in AcOH or alk. soln. The yields of the α -isomer are independent of the presence of an amino acid (pyrrolidine acid), and become greater with an increased quantity of a catalyst and a reduced time of addn. of H₂. Chas. Blanks



"APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5



APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

Census 1911

PROCESSES AND PROTECTED INDEXES

10-145-6 FM 96990

ca

10

Preparation of unsymmetric γ -glycols of the acetylene series. II.—Yu. S. Zalkind and A. I. Zakharkova. *J. Russ. Chem. (U. S. S. R.)* J., 632-3 (1933); cf. *C. A.*, 28, 23210. —See *C. A.* 27, 2079. W. P. Bricks

ASME-SEA METALLURGICAL LITERATURE CLASSIFICATION

© 2004 by the author

APPROVED FOR RELEASE: 09/19/2001 CIA-RDP86-00513R001963710009-5"

CA

The addition of hydrogen to acetylene derivatives.
 XXI. The synthesis of symmetrical diphenyl-*o*-asphaltene and its hydrogenation. Y. S. Zalkind and S. V. Nedovetskil. *J. Gen. Chem. (U.S.S.R.)*, 1, 573-7 (1931).—A warm soln. of 25 g. α -CuLiBr (II) in Et₂O added with stirring to (Bz)₂C (I), prep'd. from 6 g. Mg, 30 g. Et₂O and CuI, gave after standing 20 hrs., refluxing 8 hrs. on a Hg bath and standing 20 hrs. more, a mixt. which repd. into 2 layers. The Et₂O layer with ice H₂O and dil. AcOH yielded 1.5 g. I. The heavy lower layer by extra. with Et₂O, drying with fused K₂CO₃, distill. of the Et₂O, and recrystall. successively from PhH and EtOH, gave 17 g. (4 sym. diphenyl-*o*-asphaltene) (III) [Ph(C₆H₄C(OR)₂)₂], with 1 mol. of EtOH of crystals. Removal of the EtOH by heating at 125° for 2.5 hrs. gave II, m. 162-3°. II adds Br, does not react with KMnO₄ soln. in the cold, and gives with concd. H₂SO₄ an intense green coloration destroyed by diln. with H₂O. II (0.8 g.), heated at 140° with powdered KOH in H₂, gave I and C₆H₆, the latter identified by passing into NH₄-CuCl, forming CuS. Only 1 optical isomer of III was obtained. II is hydrogenated more slowly than [Ph(C(OH)₂)₂] (III) in EtOH with colloidal Pt, 0.005 mol. II with 10 mg. Pt requiring 8 hrs. for addn. of 2 H, 0.005 mol. III, 1.5 hrs. After addn. of 2 H, II is much more slowly absorbed. II gives 2 geometric isomers, [Ph(CuH)₂]C(OH)₂CH₂, 96% of *o*-, 4% of *o*-asphalt-1,4-diphenylhexadienol (IV) and 4% of the *o*-compd. (V). IV is difficultly sol. in EtOH and crystallizes out during the hydrogenation. IV seps. from EtOH with 1 mol. EtOH of crystals, m. 122-4° (decompn.). The EtOH-free IV thus formed or crysd. from petr. ether, m. 167°. IV adds Br more slowly than II, does not react with KMnO₄ in the cold and gives with concd. H₂SO₄ an intense green coloration changing rapidly to red and destroyed by diln. with H₂O. Oxidation of IV with KMnO₄ in AcEt with or without heating gave only CO₂. IV treated with CrO₃ in glacial AcOH gave an unidentified substance, yellow crystals, m. 163-3°. IV was the *cis*-compd. since it gave 2,6-di-*o*-asphalt-2,3-diphenyl-3,5-dihydrofuran (VI) when warmed for 1 hr. with 10 cc. Ac₂O and fused AcONa. VI was isolated from the reaction mixture by pptg. the oil by addn. of H₂O, washing with H₂O, dissolving in PhH, and pptg. with petr. ether, crystals, m. 221-2° from EtOH. VI reacts with Br but not with cold KMnO₄. IV (1.3 g.) in AcEt with 1.8 g. Pt added 2 H after 8.5 hrs. giving 2,6-di-*o*-asphalt-1,4-diphenylhexadienol (VII), m. 220°, from PhH. Addn. of 4 vol. H₂O to the mother liquor from IV gave a ppt. of IV + V. Successive recryst. from PhH and EtOH gave V, m. 231-2°, which in contrast to IV, did not crystallize with EtOH. V resembles IV in its behavior toward Br, KMnO₄ and concd. H₂SO₄. The slower the hydrogenation of II, the greater the yield of V. V (1 g. in 15 cc. AcOH) with II and 0.7 g. Pt gave VII in 2 hrs. V is the *trans*-isomer. The solubilities in 100 cc. PhH (temp. not given) were II 2.13 g., IV 0.36 g., V 0.97 g., VII 0.40 g., VI undet., but very ad. VII was also prep'd. from II in Et₂O by addn. of H₂ in the presence of Pt black.

Lewis W. Baetz

APPENDIX METALLURGICAL LITERATURE CLASSIFICATION

1930-1939

1930-1939 ONLY

CLASSIFICATION

1930-1939
1930-1939 ONLY

卷八

ZALKIND, Y.U.S.


Action of bromine on α -naphthalene sulfonic acid. Yu. S. Zalkind, L. Veisbrut and M. Alekseeva. *J. Gen. Chem. (U. S. S. R.)* 3, 802-5 (1933).—During the action of Br on α -C₁₀H₇SO₃H there are formed 3 products: 5,1-C₆H₄BrSO₃H and 1,4- and 1,5-C₆H₃Br. R. S.


-10-


ASM-SEA METALLURGICAL LITERATURE CLASSIFICATION


APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"









Esters of phosphorous acid. Yu. S. Zalikhud and E. G. Dmitrieva. Russ. 34,548, Feb. 28, 1934. Al alkylates are treated with PCl₅.

Phenol esters of phosphorous acid. Yu. S. Zelikin and R. O. Dvortseva. Russ. 34,888, Feb. 25, 1914. Phenols are heated with PCl_5 in the presence of MgCl_2 .

(P)

10

卷之三


卷之三

卷之三


METALLURGICAL LITERATURE CLASSIFICATION


卷之三

APPROVED FOR RELEASE: 09/19/2001 CIA-RDP86-00513R001963710009-5"



Methyl(chlorophenyl)carbinol. Yu. S. Zal'kind, M. K. Anusin and I. P. Berkovich. Russ. 34(630), Sept. 30, 1914. C₁₁H₁₁ClOCH₃ is hydrolyzed by heating with alkali carbonate or the like.





Tetrachlorophthalic acid. Yu. S. Zal'kina, M. V.
Belikova and O. N. Simonova. Russ. 30,701, Nov. 30,
1934. Phthalic anhydride is chlorinated in the presence
of Fe as catalyst.

CA

10

The synthesis of styrene. V. N. Zalkind, I. P. Berkovich and M. K. Amuron. *Plasticheskie Massy* 1954, No. 1, 14-20. During 4-8 hrs. KCl is added to a 3-fold excess of $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$, freshly prepnd. AlCl_3 at 70° . After boiling 1 hr. more, 75% of PhCl is obtained. The more highly ethylated compounds which are also formed may be converted to PhCl by boiling with double their wt. of $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$ and 10% of AlCl_3 . Into the PhCl contg. 1% of PCl_3 is passed Cl_2 at 70° until 80% of the theoretical amt. $\text{C}_6\text{H}_5\text{CH}(\text{OH})\text{Me}$ (I) is obtained. When boiled for 20-30 min. with a small excess of quinoline, I gives 75-80% of styrene. A cheaper process is the hydrolysis of I by boiling it with a 25% excess of a 10-15% soda soln. This gives 90% of $\text{PhCH}(\text{OH})\text{Me}$, which when dehydrated by heating with 3-5% of KHSO_4 yields 75% of styrene. App. for this

1ST AND 2ND ORDERS
3RD AND 4TH ORDERS
5TH AND 6TH ORDERS
7TH AND 8TH ORDERS
PROCESSES AND PROPERTIES INDEX

10

C.P.

Syntheses by means of vinylacetylene. Yu. S. Zel'kind and V. M. Pletz. *Trans. Leningrad. Chem.-Tech. Inst.* 1, 57-60 (1934).—The Grignard reagent from 12 g. Mg and 55 g. EtBr cooled in a CaCl_2 -snow mixt. reacted in 4 hrs. with 25 g. vinylacetylene in 75 cc. dry ether to give a sol. product vinylacetylenylmagnesium bromide (I). Twenty g. acetone was added, the reaction flask allowed to warm, and finally heated on a steam bath. After decompsn. with dil. H_2SO_4 , washing, drying over potash and (last) in vacuo, 30-3g. (32% of theory) of 2-methyl- δ -hexen-3- α -2-ol (II), $\text{H}_2\text{C}(\text{CH}_3)\text{CC(OH)Me}$, b. 75°, bp. 64° resulted. II is a yellow liquid, insol. in water, sol. in org. solvents, absorbs Br₂ and darkens with H_2SO_4 . Addn. of 20 g. of dry AcOEt to I, stirring for 2 hrs., decompsn. with ice-cold dil. H_2SO_4 in N_2 , drying over CaCl_2 and distg. at 8 mm., at 110° gave 5 g. of 3-methyl- α : β -dien-3- β -dim-5-ol (III), $\text{HOCMe}(\text{C}=\text{CHCH}_2)_2$. The mol. wt. in benzene was 143. On heating at 25 mm. III decomposes and at 1 atm. it explodes. III is insol. in H_2O , absorbs Br₂ to give an oily bromide, and at 50° reacts with anhyd. citric acid to give an orange-red hydrocarbon. Also in *J. Gen. Chem. (U. S. S. R.)* 4, 1038-90 (1934).

F. H. Rathmann

A58-51A - METALLURGICAL LITERATURES CLASSIFICATION

ITEM NUMBER	1ST 2ND 3RD 4TH 5TH 6TH 7TH 8TH 9TH 10TH 11TH 12TH 13TH 14TH 15TH 16TH 17TH 18TH 19TH 20TH 21TH 22TH 23TH 24TH 25TH 26TH 27TH 28TH 29TH 30TH 31TH 32TH 33TH 34TH 35TH 36TH 37TH 38TH 39TH 40TH 41TH 42TH 43TH 44TH 45TH 46TH 47TH 48TH 49TH 50TH 51TH 52TH 53TH 54TH 55TH 56TH 57TH 58TH 59TH 60TH 61TH 62TH 63TH 64TH 65TH 66TH 67TH 68TH 69TH 70TH 71TH 72TH 73TH 74TH 75TH 76TH 77TH 78TH 79TH 80TH 81TH 82TH 83TH 84TH 85TH 86TH 87TH 88TH 89TH 90TH 91TH 92TH 93TH 94TH 95TH 96TH 97TH 98TH 99TH 100TH 101TH 102TH 103TH 104TH 105TH 106TH 107TH 108TH 109TH 110TH 111TH 112TH 113TH 114TH 115TH 116TH 117TH 118TH 119TH 120TH 121TH 122TH 123TH 124TH 125TH 126TH 127TH 128TH 129TH 130TH 131TH 132TH 133TH 134TH 135TH 136TH 137TH 138TH 139TH 140TH 141TH 142TH 143TH 144TH 145TH 146TH 147TH 148TH 149TH 150TH 151TH 152TH 153TH 154TH 155TH 156TH 157TH 158TH 159TH 160TH 161TH 162TH 163TH 164TH 165TH 166TH 167TH 168TH 169TH 170TH 171TH 172TH 173TH 174TH 175TH 176TH 177TH 178TH 179TH 180TH 181TH 182TH 183TH 184TH 185TH 186TH 187TH 188TH 189TH 190TH 191TH 192TH 193TH 194TH 195TH 196TH 197TH 198TH 199TH 200TH 201TH 202TH 203TH 204TH 205TH 206TH 207TH 208TH 209TH 210TH 211TH 212TH 213TH 214TH 215TH 216TH 217TH 218TH 219TH 220TH 221TH 222TH 223TH 224TH 225TH 226TH 227TH 228TH 229TH 229TH 230TH 231TH 232TH 233TH 234TH 235TH 236TH 237TH 238TH 239TH 240TH 241TH 242TH 243TH 244TH 245TH 246TH 247TH 248TH 249TH 250TH 251TH 252TH 253TH 254TH 255TH 256TH 257TH 258TH 259TH 259TH 260TH 261TH 262TH 263TH 264TH 265TH 266TH 267TH 268TH 269TH 269TH 270TH 271TH 272TH 273TH 274TH 275TH 276TH 277TH 278TH 279TH 279TH 280TH 281TH 282TH 283TH 284TH 285TH 286TH 287TH 288TH 289TH 289TH 290TH 291TH 292TH 293TH 294TH 295TH 296TH 297TH 298TH 299TH 299TH 300TH 301TH 302TH 303TH 304TH 305TH 306TH 307TH 308TH 309TH 309TH 310TH 311TH 312TH 313TH 314TH 315TH 316TH 317TH 317TH 318TH 319TH 319TH 320TH 321TH 322TH 323TH 324TH 325TH 326TH 327TH 328TH 329TH 329TH 330TH 331TH 332TH 333TH 334TH 335TH 336TH 337TH 338TH 339TH 339TH 340TH 341TH 342TH 343TH 344TH 345TH 346TH 347TH 348TH 349TH 349TH 350TH 351TH 352TH 353TH 354TH 355TH 356TH 357TH 358TH 359TH 359TH 360TH 361TH 362TH 363TH 364TH 365TH 366TH 367TH 368TH 369TH 369TH 370TH 371TH 372TH 373TH 374TH 375TH 376TH 377TH 378TH 378TH 379TH 380TH 381TH 382TH 383TH 384TH 385TH 386TH 387TH 387TH 388TH 389TH 389TH 390TH 391TH 392TH 393TH 394TH 395TH 396TH 397TH 398TH 399TH 399TH 400TH 401TH 402TH 403TH 404TH 405TH 406TH 407TH 408TH 409TH 409TH 410TH 411TH 412TH 413TH 414TH 415TH 416TH 416TH 417TH 418TH 419TH 419TH 420TH 421TH 422TH 423TH 424TH 425TH 426TH 427TH 428TH 429TH 429TH 430TH 431TH 432TH 433TH 434TH 435TH 436TH 437TH 438TH 439TH 439TH 440TH 441TH 442TH 443TH 444TH 445TH 446TH 447TH 448TH 449TH 449TH 450TH 451TH 452TH 453TH 454TH 455TH 456TH 457TH 458TH 459TH 459TH 460TH 461TH 462TH 463TH 464TH 465TH 466TH 467TH 468TH 469TH 469TH 470TH 471TH 472TH 473TH 474TH 475TH 476TH 477TH 478TH 478TH 479TH 480TH 481TH 482TH 483TH 484TH 485TH 486TH 487TH 488TH 489TH 489TH 490TH 491TH 492TH 493TH 494TH 495TH 496TH 497TH 498TH 498TH 499TH 499TH 500TH 501TH 502TH 503TH 504TH 505TH 506TH 507TH 508TH 509TH 509TH 510TH 511TH 512TH 513TH 514TH 515TH 516TH 517TH 518TH 519TH 519TH 520TH 521TH 522TH 523TH 524TH 525TH 526TH 527TH 528TH 529TH 529TH 530TH 531TH 532TH 533TH 534TH 535TH 536TH 537TH 538TH 539TH 539TH 540TH 541TH 542TH 543TH 544TH 545TH 546TH 547TH 548TH 549TH 549TH 550TH 551TH 552TH 553TH 554TH 555TH 556TH 557TH 558TH 559TH 559TH 560TH 561TH 562TH 563TH 564TH 565TH 566TH 567TH 568TH 569TH 569TH 570TH 571TH 572TH 573TH 574TH 575TH 576TH 577TH 578TH 578TH 579TH 580TH 581TH 582TH 583TH 584TH 585TH 586TH 587TH 588TH 589TH 589TH 590TH 591TH 592TH 593TH 594TH 595TH 596TH 597TH 597TH 598TH 599TH 599TH 600TH 601TH 602TH 603TH 604TH 605TH 606TH 607TH 608TH 609TH 609TH 610TH 611TH 612TH 613TH 614TH 615TH 616TH 617TH 618TH 619TH 619TH 620TH 621TH 622TH 623TH 624TH 625TH 626TH 627TH 628TH 629TH 629TH 630TH 631TH 632TH 633TH 634TH 635TH 636TH 637TH 638TH 639TH 639TH 640TH 641TH 642TH 643TH 644TH 645TH 646TH 647TH 648TH 649TH 649TH 650TH 651TH 652TH 653TH 654TH 655TH 656TH 657TH 658TH 659TH 659TH 660TH 661TH 662TH 663TH 664TH 665TH 666TH 667TH 668TH 669TH 669TH 670TH 671TH 672TH 673TH 674TH 675TH 676TH 677TH 678TH 678TH 679TH 680TH 681TH 682TH 683TH 684TH 685TH 686TH 687TH 688TH 689TH 689TH 690TH 691TH 692TH 693TH 694TH 695TH 696TH 697TH 697TH 698TH 699TH 699TH 700TH 701TH 702TH 703TH 704TH 705TH 706TH 707TH 708TH 709TH 709TH 710TH 711TH 712TH 713TH 714TH 715TH 716TH 717TH 718TH 719TH 719TH 720TH 721TH 722TH 723TH 724TH 725TH 726TH 727TH 728TH 729TH 729TH 730TH 731TH 732TH 733TH 734TH 735TH 736TH 737TH 738TH 739TH 739TH 740TH 741TH 742TH 743TH 744TH 745TH 746TH 747TH 748TH 749TH 749TH 750TH 751TH 752TH 753TH 754TH 755TH 756TH 757TH 758TH 759TH 759TH 760TH 761TH 762TH 763TH 764TH 765TH 766TH 767TH 768TH 769TH 769TH 770TH 771TH 772TH 773TH 774TH 775TH 776TH 777TH 778TH 778TH 779TH 779TH 780TH 781TH 782TH 783TH 784TH 785TH 786TH 787TH 788TH 789TH 789TH 790TH 791TH 792TH 793TH 794TH 795TH 796TH 797TH 797TH 798TH 799TH 799TH 800TH 801TH 802TH 803TH 804TH 805TH 806TH 807TH 808TH 809TH 809TH 810TH 811TH 812TH 813TH 814TH 815TH 816TH 817TH 818TH 819TH 819TH 820TH 821TH 822TH 823TH 824TH 825TH 826TH 827TH 828TH 829TH 829TH 830TH 831TH 832TH 833TH 834TH 835TH 836TH 837TH 838TH 839TH 839TH 840TH 841TH 842TH 843TH 844TH 845TH 846TH 847TH 848TH 849TH 849TH 850TH 851TH 852TH 853TH 854TH 855TH 856TH 857TH 858TH 859TH 859TH 860TH 861TH 862TH 863TH 864TH 865TH 866TH 867TH 868TH 869TH 869TH 870TH 871TH 872TH 873TH 874TH 875TH 876TH 877TH 878TH 878TH 879TH 879TH 880TH 881TH 882TH 883TH 884TH 885TH 886TH 887TH 888TH 889TH 889TH 890TH 891TH 892TH 893TH 894TH 895TH 896TH 897TH 897TH 898TH 899TH 899TH 900TH 901TH 902TH 903TH 904TH 905TH 906TH 907TH 908TH 909TH 909TH 910TH 911TH 912TH 913TH 914TH 915TH 916TH 917TH 918TH 919TH 919TH 920TH 921TH 922TH 923TH 924TH 925TH 926TH 927TH 928TH 929TH 929TH 930TH 931TH 932TH 933TH 934TH 935TH 936TH 937TH 938TH 939TH 939TH 940TH 941TH 942TH 943TH 944TH 945TH 946TH 947TH 948TH 949TH 949TH 950TH 951TH 952TH 953TH 954TH 955TH 956TH 957TH 958TH 959TH 959TH 960TH 961TH 962TH 963TH 964TH 965TH 966TH 967TH 968TH 969TH 969TH 970TH 971TH 972TH 973TH 974TH 975TH 976TH 977TH 978TH 978TH 979TH 979TH 980TH 981TH 982TH 983TH 984TH 985TH 986TH 987TH 988TH 989TH 989TH 990TH 991TH 992TH 993TH 994TH 995TH 996TH 997TH 997TH 998TH 999TH 999TH 1000TH 1001TH 1002TH 1003TH 1004TH 1005TH 1006TH 1007TH 1008TH 1009TH 1009TH 1010TH 1011TH 1012TH 1013TH 1014TH 1015TH 1016TH 1017TH 1018TH 1019TH 1019TH 1020TH 1021TH 1022TH 1023TH 1024TH 1025TH 1026TH 1027TH 1028TH 1029TH 1029TH 1030TH 1031TH 1032TH 1033TH 1034TH 1035TH 1036TH 1037TH 1038TH 1039TH 1039TH 1040TH 1041TH 1042TH 1043TH 1044TH 1045TH 1046TH 1047TH 1048TH 1049TH 1049TH 1050TH 1051TH 1052TH 1053TH 1054TH 1055TH 1056TH 1057TH 1058TH 1059TH 1059TH 1060TH 1061TH 1062TH 1063TH 1064TH 1065TH 1066TH 1067TH 1068TH 1069TH 1069TH 1070TH 1071TH 1072TH 1073TH 1074TH 1075TH 1076TH 1077TH 1077TH 1078TH 1079TH 1079TH 1080TH 1081TH 1082TH 1083TH 1084TH 1085TH 1086TH 1087TH 1088TH 1089TH 1089TH 1090TH 1091TH 1092TH 1093TH 1094TH 1095TH 1096TH 1097TH 1097TH 1098TH 1099TH 1099TH 1100TH 1101TH 1102TH 1103TH 1104TH 1105TH 1106TH 1107TH 1108TH 1109TH 1109TH 1110TH 1111TH 1112TH 1113TH 1114TH 1115TH 1116TH 1117TH 1118TH 1119TH 1119TH 1120TH 1121TH 1122TH 1123TH 1124TH 1125TH 1126TH 1127TH 1128TH 1129TH 1129TH 1130TH 1131TH 1132TH 1133TH 1134TH 1135TH 1136TH 1137TH 1138TH 1139TH 1139TH 1140TH 1141TH 1142TH 1143TH 1144TH 1145TH 1146TH 1147TH 1148TH 1149TH 1149TH 1150TH 1151TH 1152TH 1153TH 1154TH 1155TH 1156TH 1157TH 1158TH 1159TH 1159TH 1160TH 1161TH 1162TH 1163TH 1164TH 1165TH 1166TH 1167TH 1168TH 1169TH 1169TH 1170TH 1171TH 1172TH 1173TH 1174TH 1175TH 1176TH 1177TH 1177TH 1178TH 1179TH 1179TH 1180TH 1181TH 1182TH 1183TH 1184TH 1185TH 1186TH 1187TH 1188TH 1189TH 1189TH 1190TH 1191TH 1192TH 1193TH 1194TH 1195TH 1196TH 1197TH 1197TH 1198TH 1199TH 1199TH 1200TH 1201TH 1202TH 1203TH 1204TH 1205TH 1206TH 1207TH 1208TH 1209TH 1209TH 1210TH 1211TH 1212TH 1213TH 1214TH 1215TH 1216TH 1217TH 1218TH 1219TH 1219TH 1220TH 1221TH 1222TH 1223TH 1224TH 1225TH 1226TH 1227TH 1228TH 1229TH 1229TH 1230TH 1231TH 1232TH 1233TH 1234TH 1235TH 1236TH 1237TH 1238TH 1239TH 1239TH 1240TH 1241TH 1242TH 1243TH 1244TH 1245TH 1246TH 1247TH 1248TH 1249TH 1249TH 1250TH 1251TH 1252TH 1253TH 1254TH 1255TH 1256TH 1257TH 1258TH 1259TH 1259TH 1260TH 1261TH 1262TH 1263TH 1264TH 1265TH 1266TH 1267TH 1268TH 1269TH 1269TH 1270TH 1271TH 1272TH 1273TH 1274TH 1275TH 1276TH 1277TH 1277TH 1278TH 1279TH 1279TH 1280TH 1281TH 1282TH 1283TH 1284TH 1285TH 1286TH 1287TH 1288TH 1289TH 1289TH 1290TH 1291TH 1292TH 1293TH 1294TH 1295TH 1296TH 1297TH 1297TH 1298TH 1299TH 1299TH 1300TH 1301TH 1302TH 1303TH 1304TH 1305TH 1306TH 1307TH 1308TH 1309TH 1309TH 1310TH 1311TH 1312TH 1313TH 1314TH 1315TH 1316TH 1317TH 1318TH 1319TH 1319TH 1320TH 1321TH 1322TH 1323TH 1324TH 1325TH 1326TH 1327TH 1328TH 1329TH 1329TH 1330TH 1331TH 1332TH 1333TH 1334TH 1335TH 1336TH 1337TH 1338TH 1339TH 1339TH 1340TH 1341TH 1342TH 1343TH 1344TH 1345TH 1346TH 1347TH 1348TH 1349TH 1349TH 1350TH 1351TH 1352TH 1353TH 1354TH 1355TH 1356TH 1357TH 1358TH 1359TH 1359TH 1360TH 1361TH 1362TH 1363TH 1364TH 1365TH 1366TH 1367TH 1368TH 1369TH 1369TH 1370TH 1371TH 1372TH 1373TH 1374TH 1375TH 1376TH 1377TH 1377TH 1378TH 1379TH 1379TH 1380TH 1381TH 1382TH 1383TH 1384TH 1385TH 1386TH 1387TH 1388TH 1389TH 1389TH 1390TH 1391TH 1392TH 1393TH 1394TH 1395TH 1396TH 1397TH 1397TH 1398TH 1399TH 1399TH 1400TH 1401TH 1402TH 1403TH 1404TH 1405TH 1406TH 1407TH 1408TH 1409TH 1409TH 1410TH 1411TH 1412TH 1413TH 1414TH 1415TH 1416TH 1417TH 1418TH 1419TH 1419TH 1420TH 1421TH 1422TH 1423TH 1424TH 1425TH 1426TH 1427TH 1428TH 1429TH 1429TH 1430TH 1431TH 1432TH 1433TH 1434TH 1435TH 1436TH 1437TH 1438TH 1439TH 1439TH 1440TH 1441TH 1442TH 1443TH 1444TH 1445TH 1446TH 1447TH 1448TH 1449TH 1449TH 1450TH 1451TH 1452TH 1453TH 1454TH 1455TH 1456TH 1457TH 1458TH 1459TH 1459TH 1460TH 1461TH 1462TH 1463TH 1464TH 1465TH 1466TH 1467TH 1468TH 1469TH 1469TH 1470TH 1471TH 1472TH 1473TH 1474TH 1475TH 1476TH 1477TH 1477TH 1478TH 1479TH 1479TH 1480TH 1481TH 1482TH 1483TH 1484TH 1485TH 1486TH 1487TH 1488TH 1489TH 1489TH 1490TH 1491TH 1492TH 1493TH 1494TH 1495TH 1496TH 1497TH 1497TH 1498TH 1499TH 1499TH 1500TH 1501TH 1502TH 1503TH 1504TH 1505TH 1506TH 1507TH 1508TH 1509TH 1509TH 1510TH 1511TH 1512TH 1513TH 1514TH 1515TH 1516TH 1517TH 1518TH 1519TH 1519TH 1520TH 1521TH 1522TH 1523TH 1524TH 1525TH 1526TH 1527TH 1528TH 1529TH 1529TH 1530TH 1531TH 1532TH 1533TH 1534TH 1535TH 1536TH 1537TH 1538TH 1539TH 1539TH 1540TH 1541TH 1542TH 1543TH 1544TH 1545TH 1546TH 1547TH 1548TH 1549TH 1549TH 1550TH 1551TH 1552TH 1553TH 1554TH 1555TH 1556TH 1557TH 1558TH 1559TH 1559TH 1560TH 1561TH 1562TH 1563TH 1564TH 1565TH 1566TH 1567TH 1568TH 1569TH 1569TH 1570TH 1571TH 1572TH 1573TH 1574TH 1575TH 1576TH 1577TH 1577TH 1578TH 1579TH 1579TH 1580TH 1581TH 1582TH 1583TH 1584TH 1585TH 1586TH 1587TH 1588TH 1589TH 1589TH 1590TH 1591TH 1592TH 1593TH 1594TH 1595TH 1596TH 1597TH 1597TH 1598TH 1599TH 1599TH 1600TH 1601TH 1602TH 1603TH 1604TH 1605TH 1606TH 1607TH 1608TH 1609TH 1609TH 1610TH 1611TH 1612TH 1613TH 1614TH 1615TH 1616TH 1617TH 1618TH 1619TH 1619TH 1620TH 1621TH 1622TH 1623TH 1624TH 1625TH 1626TH 1627TH 1628TH 1629TH 1629TH 1630TH 1631TH 1632TH 1633TH 1634TH 1635TH 1636TH 1637TH 1638TH 1639TH 1639TH 1640TH 1641TH 1642TH 1643TH 1644TH 1645TH 1646TH 1647TH 1648TH 1649TH 1649TH 1650TH 1651TH 1652TH 1653TH 1654TH 1655TH 1656TH 1657TH 1658TH 1659TH 1659TH 1660TH 1661TH 1662TH 1663TH 1664TH 1665TH 1666TH 1667TH 1668TH 1669TH 1669TH 1670TH 1671TH 1672TH 1673TH 1674TH 1675TH 1676TH 1677TH 1677TH 1678TH 1679TH 1679TH 1680TH 1681TH 1682TH 1683TH 1684TH 1685TH 1686TH 1687TH 1688TH 1689TH 1689TH 1690TH 1691TH 1692TH 1693TH 1694TH 1695TH 1696TH 1697TH 1697TH 1698TH 1699TH 1699TH 1700TH 1701TH 1702TH 1703TH 1704TH 1705TH 1706TH 1707TH 1708TH 1709TH 1709TH 1710TH 1711TH 1712TH 1713TH 1714TH 1715TH 1716TH 1717TH 1718TH 1719TH 1719TH 1720TH 1721TH 1722TH 1723TH 1724TH 1725TH 1726TH 1727TH 1728TH 1729TH 1729TH 1730TH 1731TH 1732TH 1733TH 1734TH 1735TH 1736TH 1737TH 1738TH 1739TH 1739TH 1740TH 1741TH 1742TH 1743TH 1744TH 1745TH 1746TH 1747TH 1748TH 1749TH 1749TH 1750TH 1751TH 1752TH 1753TH 1754TH 1755TH 1756TH 1757TH 1758TH 1759TH

6X
10
Esters of hydrogenated phenols. Yu. S. Zal'tkind and
M. G. Gercikov. *Plasticheskis Materialy* 1934, No. 3, 37-
40.--Cyclohexanol (I), gently heated with 10% excess
AcOH and 0.3% H₂SO₄ for 0.8 hr., yields 83% of the ester.
AcOH as weak as 6% may be used. Under the same
conditions methylcyclohexanol gives 78-80% of the corre-
sponding ester. I and 80% HCOOH, allowed to stand in
the cold without a catalyst for 1 day, give 85-90% of the
formate. I and anhyd. (C₆H₅)₂CO, refluxed without a cata-
lyst, give the diester, but the monoester cannot be so
obtained. H. M. Leicester


ASH-11A METALLURGICAL LITERATURE CLASSIFICATION

SEARCHED

SEARCHED

SEARCHED

SEARCHED



CP

The nitration of β -bromocaphthalene. Yu. Zal'kind and F. Filinov. *J. Gen. Chem. (U. S. S. R.)* 6, 979-81 (1934).—The yield of β -bromocaphthalene (**I**) obtained from β -C₆H₅NH₂ by the method of Chichibabin (*C. A.* 7, 3509) can be increased to 42% as follows: Add the diazo compd. in very small portions with energetic stirring to-freshly prepnd. CuBr and HBr at 70-80°. I stirred energetically with HNO₃ (d. 1.4) at 25-35° for 8 hrs. gave 10% of a nitrodromousphthalene (**II**), m. 110-11° from hot 30% AcOH. It was necessary to sep. the 1st crystals quickly from the still warm soln., as they were quickly followed by a flocculent material m. 76-103°. Since the 3 known nitro- β -bromocaphthalenes (1,2, 3,2 and 4,2), m. 102-3°, 89-3° and 97-8°, resp., **II** must be nitrated in the 2nd ring. **II** heated with HBr, in a sealed tube gave 1,7-dibromocaphthalene, indicating that **II** is probably the 1,7-isomer. A small quantity of the 1,2-isomer was isolated from the mother liquor from **II**. Lewis W. Butz

ASA-SEA METALLURGICAL LITERATURE CLASSIFICATION

147762-64 147762 MAP Div. 005

CA

Action of bromine on β -naphthalene-sulfonic acid. Vn.
B. Zat'kina and M. V. Belikova. J. Gen. Chem. (U. S.S.R.) 6, 1211-16 (1936); cf. C. A. 30, 6130P. By the
action of Br on β -C₁₀H₇SO₃H in an equal part of ice-cold
H₂O there is formed a mixt. of unreacted C₁₀H₇SO₃H,
Br-C₁₀H₇SO₃H and equal parts of 8,2'- and 8,2'-Br-C₁₀H₇-
SO₃H.

ASR-11A METALLURGICAL LITERATURE CLASSIFICATION

140987-14

110003 MAY 1941

RECEIVED

140987-14

Antiseptics of phenanthrores series. Yu. S. Zalkind and I. Vn. Chaitkin, Russ. 40,148, Jan. 31, 1937. Phenanthrores are heated with an excess of $(\text{NH}_4)_2\text{SO}_4$ and NH_3 to 140-150°; phenanthrylamine are sepd. from the reaction mixt. by soln. in acid and a reppin. with alkali.

CA


卷之三


כט

410.11A METALLURGICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"






Investigations in the field of vitamin A synthesis. Yu.
S. Zel'kind, S. Zosin and N. Blokhin. *Comp. rend. Acad.*
U. R. S. S. R., 57-60 (in German 61-3) (1933).— β -
Iucone and the Mg deriv. of vinylacetylene were heated on
a H₂O bath for 2 days till the odor of iucone disappeared.
After this usual treatment a viscous liquid, β -methyl- β -
(γ , δ , δ' -trimethyl- γ -butyryloxy)- β -hydroxy- α -methyl- β -
MeC(CH₃)₂CMe:CH₂:CHCMe(OH)C:CH₂—
(I),


Cl_{fr}, bp. 135-45°, d₂₅²⁰ 1.0384, was obtained.
This was reduced by almost the caked. vol. of H₂ and a Pt
catalyst to β -(γ , δ -trimethylcyclohexyl)ethyldimethylbutylic
carbinol, MeC(CH₃)₂CMe:CH₂(CH₂)₂Me(OH)(CH₂).

Cl_{fr}, bp. 125 0°, which was identified by oxidation
with K₂Cr₂O₇. This gave tetrahydroiuanone (II) and
PrCO₂H. The latter appeared to be contaminated with a
little trimethylcyclohexylacetic acid. II was identified by
comparison with tetrahydroiuanone, bp. 112-13°, d₂₅²⁰ 0.9833,
d₂₅²⁰ 0.9224, n_D²⁰ 1.4766 (semicarbazone, m. 105-6.5°),
obtained by hydrogenating iucone with H₂ and Pt. While
I had many of the chem. characteristics of vitamin A,
tests on rats showed that it was not a vitamin.

John E. Millery

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION





XIII. Bis(1-hydroxyethylacetyl)acetylene derivatives. VII. N. Zal'kigul. *J. Russ. Chem. Soc.* (U. S. S. R.) 33, 1723-7 (1933); *cf.* *C. A.* 28, 2707. — Previous study of the rate of addition of H₂ to acetylene γ -glycols showed that secondary glycols are hydrogenated more slowly than bis-secondary glycols (*z.* and *Nedzhad, J. Russ. Phys.-Chem. Soc.* 50, 36 (1918)), and that the reaction velocity decreases with the increasing mol. wt. of the radicals (*cf. Z. and Bulyat'yakov, J. Russ. Phys.-Chem. Soc.* 47, 1830 (1915); *Z. and Markaryan, C. A.* 11, 584). The influence of cyclic radicals, connected directly with the HO-bearing C, on the rate of hydrogenation of bis-acetylene glycols was studied by hydrogenating 0.01 mol. alc. in the presence of various amounts of Paul's colloidal Pd at 18-21° and 750-810 mm. pressure, *i.e.*, by 183° (decomp.), m. 109° (103°, Dupont, *C. A.* 8, 1579), was prepnd. by the Dupont method. It gives with AcO and KOAc the diacetate, m. 45.5-6.5°, and with Br-

$\text{C}_6\text{H}_5\text{CH}_2\text{C}(\text{Br})\text{Cl}_2\text{C}_6\text{H}_5$, m., 100.5-7.5%. Like other bimary glycols, it easily adds 2 H atoms, after which the rate of reaction declines. It differs from other glycols with radicals of nearly equal or even smaller magnitude. It is more easily hydrogenated than $(\text{MeC}(\text{OH})\text{C}_2)$, (II) and $(\text{EtC}(\text{OH})\text{C}_2)_2$, (III). The rate of hydrogenation of $(\text{MeC}(\text{OH})\text{C}_2)_2$, II and III (18, 85 and 145 min., resp.) (loc. cit.), while that of I (30 min.) shows no relation to other glycols. The probable stereochem. causes influencing the reaction are being investigated. With increasing ratios of the catalyst (1, 2, 4, 8 and 10), the hydrogenation velocity at first increases at a disproportionate rate and then imperceptibly drops (1, 5.6, 20.1, 29.4 and 47.8, resp.). Under all conditions of hydrogenation only *1-butyl-1-hydroxyethylene* (ethylidene, m., 162°) is obtained. It is difficultly oxidized, giving only 0.5 g. adipic acid on oxidizing 5 g. of the glycol with 5.24 g. $\text{K}_2\text{Cr}_2\text{O}_7$ (5% soln.) at room temp. for 12 hrs. **XIII.** Dihydro-*butyldiol* and its hydrogenation. Yu. S. Zal'kind and V. I. Tretiashvili. *Ibid.* 1708-71.—The effect of substitution of an open chain for a Ph radical of an equal no. of C atoms in an acetylene given on the rate of addn. of


APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

H_2 was studied by a comparative hydrogenation of di-
methylbenzene (I) and diphenylbenzene (II) (cf. Z. and
Neubrath, loc. cit.), 1, m. 47-49°, was obtained in 97%
yield by the method of Lotrich (Z. Russ. Phys.-Chem. Soc.
18, 656 (1903)) and Dupont (loc. cit.). I after repeated,
recryst. from ligroin sepd. a few crystals of a compd., m.
68.7-70°. The mol. wt. of 1 is 211 ($\text{Me}_2\text{C}_6\text{O}$) and 210
($\text{C}_6\text{H}_5\text{C}_6\text{H}_5$) (calcd. 254.2). I was hydrogenated in the presence
of colloidal Pd as described above. The reaction
was completed in 15 hrs, requiring an addnl. introduction
of Pd after 6 hrs. of hydrogenation. A break in the rate of
hydrogenation, after the addn. of the 1st 2 H atoms is
sharply defined. With 1 mg. and 7.5 mg. of the catalyst,
the relative addn. of H for the 1st 6 min. is, resp., 12.76
and 65.10% for I, 7.8% for II and 20.44 and 59.17% for
dimethylbenzene. Thus a substitution of benzyl for
Ph, but not for a lighter radical, considerably accelerates
the hydrogenation. I on the addn. of 2 H atoms gives 2
ethylene glycols, [$\text{Me}(\text{CH}_2)\text{CH}(\text{OH})\text{CH}_2$]_n, cryst., m.
44-5°, and liquid, b.p. 186-201°, d²⁵ 0.84. The cause of
the formation of 2 ethylene glycols may be that I is a mixt.
of d + l and l-isomer, or that in the hydrogenation of I
there are formed 2 geometrical isomers corresponding to
the same acetylene glycol. The same correlation of the
reaction products obtained by the rapid hydrogenation
(8-10 min. Pd) and by the slow hydrogenation (1 hr. Pd)
disproves the 2nd postulation. Furthermore in case of
geometrical isomorphism, an exhaustive hydrogenation of the
ethylene glycols in the presence of Pt black catalyst
would give identical and, if possible, because the asym. C
atoms are not affected in hydrogenation. I on the addn.
of 4 H atoms gave 2 products, m. 105-6° and m. 65-6°,
analysing for hexadecane-1,10-diol. Exhaustive hy-
drogenation of the cryst. ethylene glycol gave only 1 said.
glycol, m. 85-6°, while that of the liquid ethylene glycol
gave 2 isomers in the ratio of 1:1. It follows that the 2
ethylene glycols are not geometrical isomers but are de-
rived from the 2 isomeric I. I treated with Br_2 in CHCl_3 ,
gave 45% of diketone, m. 97-8° (Dupont, loc. cit.).
This on cleaving Br with Zn dust in alc. gave 1, m. 13-4°,
which on hydrogenation with colloidal Pd gave only the
cryst. ethylene glycol, m. 44-5°, and with Pt black the
liquid glycol, m. 85-6°.

"APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5



APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

Production of tetrachlorophthalic acid. Yu. S. Zel'kind and M. Bedikova. *J. Applied Chem. (U.S.S.R.)* 8, 1210-13 (in German 1213) (1935).—A yield of 90-5% $\text{C}_6\text{Cl}_4(\text{CO}_2\text{H})_2 \cdot \text{H}_2\text{O}$ (I), m. 244-5° (open tube), is obtained by chlorinating $\text{C}_6\text{H}_4(\text{CO}_2\text{H})_2$, (II) in the presence of 3% Fe filings at 240-30° to the correct m. p. of the reaction product. The chlorination without stirring proceeds slowly, requiring 63-8 hrs. for 300 g., and 110 hrs. for 2 kg. II. $\text{C}_6\text{H}_4(\text{COCl})_2$ reacts with Cl more rapidly, giving 98% I. It loses the water of crystal. at 100° and the water of constitution at 110°, giving $\text{C}_6\text{Cl}_4(\text{CO})_2\text{O}$.

10

60

卷之三

卷之三

AMERICAN METALLURGICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

ca

10

o-Vinylnaphthalene and some of its derivatives. Yu. S. Zalkind and S. A. Zomis. *J. Gen. Chem. (U.S.S.R.)*, 29, 6788-681 (1959). — The polymerization ability of high-mol.-unsatd. compds. in relation to their structure and the properties of the resulting polymers was studied in the case of *o*-vinylnaphthalene (**I**) and its derivs. substituted in the side chain. **I**, RCH:CHMe (**III**), RCMe:CH₂ (**III'**), RCMe:CHMe (**IV**), and RC≡CHMe (**V**) (R = OEt, were prep'd. by the Grignard reaction by interaction of C₆H₅MgBr and the corresponding alcs. with subsequent dehydration of the resulting idcs. with KHSO₄ (cf. Shorygin and Shorygina, *C. A.*, 59, 18412). The alcs. were identified as the acid esters of C₆H₅CO₂H. The polymerization was effected without catalysts and in the presence of Br₂O₂H and floridin at 20°, 40°, 60° and at higher temps. The polymerization rate was detd. either by vacuum distn. of the monomer and weighing the polymer residue, or by measuring the relative viscosity in the Ostwald viscometer. Under all conditions gave only a polymer and in the amt. proportional to the elapsed time. The viscosity of polymerization product increased at a greater rate than the percentage of polymer in the reaction mixt. Thus, viscosities of 9.81 and 30.28 with 10.28% and 20.63% polymer resulted after 11 and 22 days, resp. The viscosity of a polymer at an equal concn. obtained at 20° is greater than that of the product obtained


at 40°. The resistance to heat is 150-180° for I polymerized at 20°, and 110-20° for the product obtained at 20°. The polymer of **I** obtained in the presence of Br₂O₂H catalyst showed poor insulating properties and considerable dielec. losses. The derivs. of **I** showed poor polymerizing abilities even at elevated temps. (boiling water bath). **V** failed to polymerize under all conditions. **III** and **IV** showed no increase in the viscosities after 44 days of polymerization in the presence of 0.5% Br₂O₂H, while **II** gave 8.2% polymer. **III** and **IV** doubled the viscosities in the presence of floridin after 100 days at 20°. The substituted **I** gave chiefly dimers and no polymers as in the case of **I**. The sym. derivs. of **I** are more easily polymerized than the asym. ones. **I**, b.p. 105-0°, d₄²⁰ 1.0774, n_D²⁰ 1.6385, M. R. 52 (60.41 calcd.), exaltation 1.60. **II**, b.p. 112-13°, d₄²⁰ 1.0429, d₄²⁵ 1.0283, n_D²⁰ 1.6335, exaltation 3.51, M. R. 55.03 (57.84 calcd.). **III**, b.p. 112-13°, d₄²⁰ 1.027, n_D²⁰ 1.6142. **IV**, b.p. 113.5-115°, d₄²⁰ 1.0383, n_D²⁰ 1.6137, exaltation 0.4, M. R. 60.03 (59.45 calcd.). **V**, b.p. 119-20°, d₄²⁰ 1.034, n_D²⁰ 1.507, exaltation 0.35, M. R. 64.02 (64.37 calcd.). More than 30 references. Chas. Blane

ASLB-SLA METALLURGICAL LITERATURE CLASSIFICATION

SEARCH STRATEGY												FILING NUMBER											
SEARCHED	INDEXED	SERIALIZED	FILED	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	
12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	

Adipic acid. Yu. S. Gal'kind and L. F. Markov. Russ. 50,301, Jan. 31, 1937. Adipic acid is prep'd. by oxidizing cyclohexene with solns. of $K_2Cr_2O_7$ in the presence of $HgSO_4$.

ABQ-SLA - METALLURGICAL LITERATURE CLASSIFICATION



Unsaturated alcohols - V. A. Zel'nik and B. A. Zel'nik
Russ. 814006, Oct. 31, 1957. Also, vinyl, at least one
triple, and two double bonds are prep'd. from acetylene
bivinylmagnesium halides by the action of aldehydes or
ketones.

Ca

10

Obtaining esters of dicarboxylic acids. Yu. S. Zel'kind and M. G. Gerzhikov. Narednyi Komissariat Tsvetkoi Prom. S. S. R., Nauch.-Issledovat. Inst. Plasticheskikh Mass., Plasticheskie Massy, Sbornik 2, 191-7(1937).—Yields of 85-95% of the following adipates are obtained by standard methods: dicyclohexyl, m. 37°; bis(methylcyclohexyl); di-Bt; diisomanyl, b.p. 190°, d_{4}^{20} 0.9455, n_{D}^{20} 1.4343; diisobutyl, b.p. 186-8°, d_{4}^{20} 0.9542, n_{D}^{20} 1.4301; di-Bu b.p. 190-1°, d_{4}^{20} 0.9550, n_{D}^{20} 1.4339. H. M. Leinweber

C

A

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

S

The catalytic dehydrogenation of borneol and isoborneol.
Yu. S. Zal't'kin, Yu. I. Mikhelis, M. A. Stoyakov and
R. L. Levin-Kogan. Naredayi Komissariat Trudovoi
Prom. S.S.R., Nauch.-Issledovat. Inst. Plasticheskikh
Mass, Plasticheskie Massy, Shevchenko 2, 243 81 (1937).
Ni(OH)₂, reduced with H₂ at 250-300° needs no activator
for liquid-phase dehydrogenation of borneol. Isoborneol
is more difficult to dehydrogenate. Cu₂(OH)₃CO₂ is a
cheaper and more stable catalyst. It should be mixed
with MgCO₃ to increase the surface. Boiling the borneol
for a short time with CaO or Ca(OH)₂ before reaction
eliminates traces of acids which slow the dehydrogenation
and makes the catalyst more active. II. M. Leicester

The synthesis of camphor by the method of J. Ritter.
Yu. S. Zal'kind and A. Rgurova. *Narodnyi Komissariat
Tsvetkov Prom. S. S. S. R., Nauch.-Issledovatel. Inst.
Plasticheskikh Mirev, Plasticheskie Mery, Sbornik 2,
121-30(1937); cf. C. A. 37, 4780.—Minor modifications
are suggested.* H. M. Leicester

10

45-1144 METALLURGICAL LITERATURE CLASSIFICATION

SEARCHED	INDEXED	SERIALIZED	FILED
1	2	3	4

1ST AND 2ND COVERS

PROCESSED AND RECEIVED

CH

Obtaining adipic acid from oil distillates. Yu. S. Zal'kina and I. P. Markov. *Narodnyi Komissariat Tsvetnoy Prom. S. S. S. R. Nauch.-Issledovatel. Inst. Plasticheskikh Mass., Plasticheskie Massy, Sbornik 2, 331-44* (1937).
The cyclohexane obtained from the 80-110° fraction of an oil distillate can be oxidized directly to adipic acid by conc. HNO₃, but the yield is low. Continuous chlorination of the cyclohexane gives 61% cyclohexyl chloride which easily splits out HCl when passed over activated C with steam at 280-300° to give 92% cyclohexene. Oxidation of the latter with a mixt. of H₂SO₄ and Na₂Cr₂O₇ gives 45-49.5% adipic acid. H. M. Leicester

10

ASW-LSA METALLURGICAL LITERATURE CLASSIFICATION

二八

A new synthesis of bicyclic glycols. Yu. S. Zel'kind and M. A. Al'tikovich. *J. Gen. Chem. (U.S.S.R.)*, **7**, 227-33 (1937); *cf.* *J. Am. Chem. Soc.*, **30**, 2323 (1908).—In a previous study of the condensation of amino-substituted acetylenes in the presence of $\text{CuCl} + \text{NH}_4\text{Cl}$ acidified with HCl , it was found that $\text{PhC}\equiv\text{CH}$ gives diphenylacetylene. The application of this reaction to the synthesis of bicyclic derivatives of OH compounds was begun with tertiary acetylenic alcohols. The reaction leads to the formation of tertiary bicyclic glycols, according to the scheme: $2\text{RC(OH)C}\equiv\text{CH} \rightarrow [\text{RRC(OH)}\text{C}(\text{C}_2\text{H}_5)\text{C}_2\text{H}_5]_2 + \text{H}_2$. Thus, $\text{MeC(OH)C}\equiv\text{CH}$ (**I**) gave $[\text{Me}_2\text{C(OH)}\text{C}(\text{C}_2\text{H}_5)\text{C}_2\text{H}_5]_2$ (**II**) and 1-hydroxybicyclohexylacetylene (**III**) gave bis(1-hydroxybicyclohexyl)acetylene (**IV**). **I**, b. 103.5-4°, $d_4^{20} 0.8832$, n_D²⁰ 1.4202, M. R. 24.08 (calcd. 24.81) (*cf.* Pavorski, U. S. S. R. pat. 31,017), **II** was introduced into 2 g. CuCl and 6 g. NH_4Cl in 25 cc. H_2O at room temp. The reaction was completed in 3-5 min., giving 100% **II**, m. 132-2.8° (CaH_2). **II** in BzOAc hydrogenated in the presence of Pt black added 8 H atoms, forming 2,2-dimethyloctane-2,7-diol, m. 48-9° (lignin), 57-8° (H_2O) (*cf.* Zel'kind, *J. Russ. Phys.-Chem. Soc.*, **38**, 931 (1906)). The reaction proceeds without the usual addition of HCl and gives equally good results in the absence of NH_4Cl and with NH_4OH soln. of CuO . Since the reaction affords 100% **II** without the participation of Cu and evolution of a gas, it might be assumed that the atm. C_2 acts as an acceptor of the H_+ .

liberated in the reaction. In fact, the reaction in a H atom, proceeds very slowly, giving after 1 hr. but few crystals of **II**. In an atm. of pure O₂, the condensation takes place very rapidly with a decrease of the O₂ vol., corresponding to the atm. required for oxidizing **Cu(O)** and the H₂ liberated in the reaction. **III**, b.p. 99-101°, d₂₀²⁰ 0.9058, n_D²⁰ 1.4750, M. R. 30.17 (calcd. 30.47), obtained by the Rupe method (C. A. 22, 2929), was contaminated with about 8% cyclohexanol. **IV**, I, m. 173-4° (CuI), resulted in nearly 100% yield from 3.1 g. **III** and 3 g. CuCl and 9 g. NH₄Cl in 30 cc. H₂O. The reaction is completed at room temp. in 20-3 min. in O₂ and in 2-3 hrs. in air. In N an insignificant amt. of **IV** resulted after 12 hrs. of interaction. The condensation of **III** in the presence of CuO in NH₄OH gave 0.2-3% **IV**. Hydrogenation of 2 g. **IV** with 2 g. Pt black resulted in the addn. of 1/2 H atoms and formation of *α,β*-dihydroxybutane, b.p. 109°, and with 0.6 g. Pt black in冰水, dihydrocyclohexyl(1,4-butane) Chas. Blame.

10

PROCESSES AND PROPERTIES INDEX

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION

卷之三

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

CROSS REFERENCE

OPEN

CLOSE

MATERIALS

PAPER

- cc*
1. The addition of hydrogen to acetylene compounds. XXVII. The catalytic hydrogenation of 2,5-dimethyl-1,3-hexadien-3-one. Yu. S. Zel'kind and Z. V. Sesagina. *J. Gen. Chem. (U.S.S.R.)*, 7, 470-5 (1937); cf. *J. A. 31*, 1012^a.—This compd. is best prepd. by dehydration of the corresponding glycol with H_2SO_4 . When it is hydrogenated over Pt, the normal stdn. hydrocarbon is produced, but if Pd is used, only 6 H atoms are absorbed. The first step is probably to $CH_3:CHMeCH_2CHMe:CH_3$.
2. This then gives chiefly 2,5-dimethyl-1-hexene, though the product seems to contain not more than 3-5% of 2,5-dimethyl-3-hexene. H. M. Leicester

10

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION									
SUBJECTIVE		OBJECTIVE							
SEARCHED		INDEXED							
SEARCHED		INDEXED							
SEARCHED	INDEXED	SEARCHED	INDEXED	SEARCHED	INDEXED	SEARCHED	INDEXED	SEARCHED	INDEXED


The addition of hydrogen to acetylene derivatives
XVII. Hydrogenation of symmetrical diphenyltolyl-
butinediol. Yu. S. Zalkind and E. R. Martinson. *J.
Am. Chem. C. S. S. R.* 7, 815-17 (1937); cf. *C. A.* 31,
1404. — Continuing the study on the hydrogenation of


tetraarylbutinediols to give both possible geometric isomers of the olefin glycid, *1,4-diphenyl-1,4-di-p-tolylbutene-1,6-diol* (**I**) is synthesized and its behavior on hydrogenation shown to conform with that of previously studied tetraarylbutinediols. To $\text{BnMgCl} \cdot \text{C}_6\text{H}_5\text{Cl}$, prep'd. from 8 g. Mg and 35 g. Et_2Br , is added 80 g. (15 g. excess) $p\text{-MeC}_6\text{H}_4\text{COPh}$ (**IV**), m. 57° , prep'd. by the Friedel-Crafts reaction from BnCl and PhMgBr , the reaction mixture allowed to stand 5 days, decomposed with dil. AcOH , and the ppt. (48 g.) recrystd. twice from EtOH to give 25 g. (35.0%) **I**, m. 140° . The structure of **I** is proved by fusion with KOH in a stream of H_2 to yield **CII** and **IV**, m. 57° . **I** (2.09 g.) in dry Et_2O contg. Pt black (0.3 g.) is completely hydrogenated in 90 min. to *1,6-diphenyl-1,6-dip-tolylbutene-1,6-diol* (**II**), crystals from acetone, m. 176° . **I** (2.09 g.) in 100 ml. EtOH contg. colloidal Pt (10-50 mg.), hydrogenates with some difficulty to give the 2 geometric isomers of *1,4-diphenyl-1,4-di-p-tolylbutene-1,6-diol* (**III**): (trans form (10.8% yield), crystals from acetone or alc., m. 188.90° ; cis form (70.7% yield), crystals from ligroin, m. 90°). Both forms of **III**, in the presence of Pt black, reduce to **II**. Concentrated H_2SO_4 is an excellent reagent to distinguish between **II** and the 2 forms of **III**: **II** gives a green-yellow coloration, while both forms of **III** give a characteristic crimson color, passing rapidly through a series of intermediate colors to a red-green. J. L.

ABE-31A METALLURGICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"





The addition of hydrogen to acetylene derivatives.
XXIX. The action of sodium and alcohol on acetylene glycol and the geometric isomers of tetramethylbutenediol.
 Yu. S. Zal'kind and S. V. Lukhovets. *J. Gen. Chem. (U.S.S.R.)* 7, 2417-22(1937); cf. *C. A.* 31, 20281.
 Bourquel and Rambaud (*C. A.* 34, 2720) claim to have obtained a *trans*-tetramethylbutenediol, m. 101-2° (I), from tetramethylbuteniodiol (II) by reduction with Na and EtOH, and that the compd. of Z. m. 75-6° (III) is an impure mist. When II is reduced over Pt it adds H atoms and gives tetramethylbutanediol. Over Pd it adds 2 H atoms to form III. When it is boiled with Na and iso-AmOH it forms C_2H_2 , a very sensitive test for acetylene alcs. It forms the same dibromide that is obtained from II. It is therefore a dimorphic form of II. When melted II is seeded with I, the solid product is mostly I though the reverse change does not occur. III is actually the *trans*-tetramethylbutenediol.

4.3.3.1.4. METALLURGICAL LITERATURE CLASSIFICATION

100-43413-X

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

Catalytic oxidation of phenanthrenes by air. Vg. 9.
Zel'kind and V. V. Kecarev. *J. Applied Chem. (U.S.S.R.)* 10, 99-104 (in German) 104 (1957).—The oxidation of phenanthrene in the presence of humic impregnated

with 36% V_2O_5 at 400° (1 hr. temp. of the phenanthrene was 275°, which corresponds to 3% of phenanthrene in the gas mixt.) by O_2 and by air to which 12% of O_2 was added, yielded CO_2 , H_2O and only 2% of phthalic acid. Preheating phenanthrene to 225° and carrying the reaction out at 370° and a duration of contact of 4 sec., the yield of phthalic acid was highest (11.5%). Added CO_2 to the air did not increase the yield of phthalic acid, but increased moisture content of the air yielded 14% of solid acids in 177°, contg. mainly phthalic acid and some diphenic acid (in. 210°); further increase of the moisture to 60% did not improve the yield. The best yield (21%) of acids was obtained by using paraffine with a concn. of NH_4VO_3 equal to 25%. The gradual introduction of phenanthrene also increased the yield of acids; this can be explained by gradual increase of the activity of the catalyst. Pure WO_3 did not catalyze the reaction at all, and its mixt. with V (W 20 and V 80%) yielded only 12% of acids. The mixt. of Mo (II) and V 90% catalyzed the reaction, yielding 30% of acids, but pure Mo is a poor catalyst. The mixt. of U_2O_5 and V_2O_5 (U 30 and V 70%) catalyzed the reaction to a greater extent, yielding 41%

The optimal conditions for the last catalyst were 10 g. of the catalyst, a contact time of 1.2 sec., reaction temp. of 430° (if the phenanthrene was heated to 240°) or 440° (if it was heated to 235°). The last catalyst was very stable; after 12 expts. its activity had not decreased at all. The catalyst composed of V 10, Mo 10 and U 30%, at 330-350° (the phenanthrene was heated to 218°) and a contact time of 2 sec., yielded 30% of solid acids. Addn. of the oxides of Zn, Cu, Ag, Bi in the U-V catalyst had practically no influence on the yield. The acids obtained had in all cases a yellow color, m. 160°, and consisted of phthalic acid, which partially changed to BaOII; in some expts. a small yield of diphenic acid and phenanthrenequinone was obtained, and, in some, the acid products treated with H₂SO₄ gave a red color, characteristic of fluoresceinophthalic acid, and the neutral products treated with concd. H₂SO₄ gave a violet color, characteristic of fluoresceone. The oxidation of phenanthrene was also carried out in liquid phase, by passing a stream of the air or O₂ through the liquid phenanthrene (at 240°), but the yield was very poor. In this case, the oxidation with air yielded diphenic acid, and that with O₂ maleic acid. The mechanism of the oxidation of phenanthrene is given as follows: phenanthrene → phenanthrenequinone → 6,6'-diphenic acid → *s*-phthalic acid → BaOII → maleic acid → CO + H₂O; and the formation of the fluoresceone and fluoresceinophthalic acid can be explained by the following scheme: 6,6'-diphenic acid → 2,6'-diphenic acid → fluoresceinophthalic acid → fluoresceone → *s*-phthalic acid. Four references.

A. A. Podgorny

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

Preparation of the metacrylic acid ester from isobutylene
acid. Yu. S. Zal'kind and I. P. Markov. *J. Applied
Chem. (U.S.S.R.)* 10, 1022-4 (in German 1044) (1937).—
Isobutylene acid was chlorinated in the presence of elec-
tric light (20-w. globe), starting the reaction at first at 50°;
then the temp. could be dropped to 20-30° without
diminishing the velocity of the reaction. The product
was fractionated, yielding unchanged acid, α - and β -
chloroisobutyric acid. Under the above conditions the
yield of β -isomer was the greatest. The Me esters of
both isomers were prep'd. by heating the acids with 2
parts of MeOH in the presence of H₂SO₄ (conc'd.) for 3
hrs. The resulting mixts. were neutralized with soda
(aq.) and the ester of the α -isomer was extd. with Et₂O,
whereas the ester of the β -isomer was collected in the
lower layer and washed with water to neutral reaction.
The yield was about 70% (theory). The ester of the
 α -isomer bp 64-65°, bnd 126-0.5°, d₄²⁰ 1.0033, n_D²⁰ 1.4019,
M. R. 31.82; the β -isomer bp 85-90°, bnd 151-5°, d₄²⁰
1.1101, n_D²⁰ 1.4297, M. R. 31.60. The splitting off of
HCl from the esters was accomplished by boiling with
1.5 parts (by wt.) of quinoline in the presence of a small
amt. of hydroquinone (depolymerization agent). The
yield for the α -isomer was 68% and for the β -isomer
87.4%; and the reaction proceeded twice as fast in the lat-

ter case as in the former. The product had all the proper-
ties of the Me ester of metacrylic acid. The splitting
off of HCl from α - and β -chloroisobutyric acids was accom-
plished by boiling with 2% of activated vegetable char-
coal in the presence of a small amt. of hydroquinone.
The product of the reaction was distd. at 165° and had
all the properties of metacrylic acid, which was esterified
with MeOH. Seven references. A. A. Podgorny

ASB-SEA METALLURGICAL LITERATURE CLASSIFICATION

STANDARD SUBJECTS

SUBJECT WORDS OR KEY WORDS

The addition of hydrogen to acetylene derivatives. XXX. Catalytic hydrogenation of symmetrical dimethyl-disopropylbutynediol. Yu. S. Zel'kind and A. I. Nogafdei. *J. Gen. Chem. (U. S. S. R.)* 8, 1382 (1938); *J. C. d.* 32, 2880. To $(\text{CH}_2)_2\text{CO}_2$, prep'd. from 137 g. Et_2Br and 32 g. Mg in 200 ml. Et_2O , is added dropwise 121 g. $\text{Me}_3\text{CCHCOMe}$ at 0° in 3 hrs.; the reaction mixt. allowed to stand 12 hrs., decomposed with dil. AcOH , extd. with Et_2O and the residue from Et_2O is fractionated to give $[\text{CC}(\text{OH})(\text{CH}_2\text{Me})_2]_2$ (I), b.p. 120-2°, m. 78-80° (digruin). I in alc. contg. colloid Pd is hydrogenated to *s*-dimethyl*dis*propylbutenediol, b.p. 110-21°, d₄²⁰ 0.9344, n_D²⁰ 1.4605. I is hydrogenated 3 times slower than dimethylidethylbutynediol but considerably faster than tetracylbutynediol. Chas. Blanc

Action of bromine upon a glyned of the ethylene series
 Vito, N., Zalkind and A. L. Nageloff, J. Am. Chem. Soc., 61, 1810 (1939); cf. Zalkind and Peterlin, C. A., 30, 2077.—[PbCH(OH)CH₃]_n (I) (m.p. 152°) in CHCl₃, on treatment with Br at 0° gives 2,5-diphenyl-3,6-dibromohexa-2,5-dienoic acid (II) and 2,5-diphenyl-3-bromo-2,5-dihydrofuran (III). The reaction at -15° gives only II isolated by direct vacuum distn. of the solvent at room temp. It gradually, though incompletely, loses HBr to form III. Neither bromide gives CH₃ with LiAlH₄. I adds I even more sluggishly to form the monoadduct. The reaction with [iso-PrMg(OH)Cl] gives 2,5-dimethyl-2,5-dipropyl-3-bromo-2,5-dihydrofuran, d₄ 1.2942. A mixt. (9 g.) of stereoisomeric dimethylbenzylidene-3-hexene-2,5-dinols (IV), b.p. 104-11° and 124-4 g. Br in 12.0 ml. CHCl₃ gave 0.7 g. 3,6-dibromo-2,5-dioxane diol, m.p. 139.5-40°, and 14.5 g. (100%) 2,5-dimethyl-3,6-dibromohexa-2,5-dienoic acid. When vacuum distd., it cleaves 188° to give 2,5-dimethyl-3-bromo-2,5-dihydrofuran, b.p. 48-51°. An ester of IV, prepared by hydrogenation of benzylidene diol acetate (cf. Z. and Bezonova, C. A., 30, 2327), gave 2 fractions, b.p. 80-7° and 84-8°. The fraction, b.p. 80-7°, adds 2 H atoms in the presence of the Pd catalyst to form 2,5-hexanediol, b.p. 83-4°, d₄ 0.9983, n_D²⁰ 1.4232; M. R. n 31.02. The fraction, b.p. 84-8°, reacts with Br to give the monobromoglycidyl, MeC(CH₂OCH₂)₂CH₂Br(IV), b.p. 92-10°.

ASM-SEA METALLURGICAL LITERATURE CLASSIFICATION

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

CQ

10


Structure of several polychloro derivatives of biphenyl.
 Yu. S. Zalkind, and M. V. Belikova. *J. Gen. Chem. (U.S.S.R.)* 8, 1918-21 (in French). (1921) (1938).
 KMnO₄, CrO₃ and HNO₃ (d. 1.2) have very little or no
 orientating effect on polychlorobiphenyls. Pentachloro-
 biphenyl (I), bp 195-220°, d₂ 1.6, when oxidized with
 HNO₃ (d. 1.4) for 100 hrs. gives a mixt. of 3,4-Cl₂C₆H₄-
 CO₂H (II), m. 204-5°, and 2,4,5-Cl₃C₆H₃CO₂H (III), m.
 190-90.5°. Heptachlorobiphenyl (IV), bp 240-280°, d₂
 1.64, when oxidized gives a small amt. of III and 2,3,4,5-
 Cl₄C₆HCO₂H (V), m. 190.5-1.5°. To confirm its struc-
 ture V was prep'd. by heating o-(CO₂H)₂C₆H₄ in AcOH to
 300° for 7 hrs.. I is therefore the 2,4,5,3',4'-compd. and
 IV is 2,3,4,5,2',4',5'-heptachlorobiphenyl. The benzoic
 acids were isolated as the amides: amide of II, m. 129-
 30°; III, m. 107-7.5°; V, m. 190-2°. In Ph, the Cl
 evidently has a greater orientating influence than does Ph.
 John Livak.

Some derivatives of cyclopentane. Yu. S. Zal'kind and I. F. Markov. *J. Applied Chem.* (U. S. S. R.) 11, 818-22 (in French 822) (1938); cf. *C. A.* 31, 28751.—The chlorination of cyclopentane at room temp. yielded 75.3-79.7% (on reacted hydrocarbon) of cyclopentyl chloride which easily split out HCl when passed over activated C with steam at 350-70° to give 88% (on reacted chloride) of cyclopentene. Treatment of the latter with 80% soda at 22-38° yielded 81.2-82.2% cyclopentane. Boiling the mixt. of adipic acid with a double amt. of cyclopentane in the presence of 1% by wt. of the adipic acid used of concd. H_2SO_4 , under a reflux condenser for 4 hrs., yielded 89.9% of dicyclopentyl adipate, m. 31-3°, η_{D}^{20} 262-54°. Similar treatment of cyclopentane with glutaric acid yielded 89.4% of dicyclopentyl glutarate, m. 218-20°, d_{10}^{20} 1.0392. Similar treatment of cyclohexanone with glutaric acid yielded about 100% dicyclohexyl glutarate, m. 236-8°, d_{10}^{20} 1.0415. Both esters had better plastic properties on nitrocellulose than dicyclohexyl adipate or trityl phosphate. On benzylcellulose both esters are about equal in their plastic properties to dicyclohexyl adipate and higher than trityl phosphate. Fifteen references. A. A. Podgorny.

After review
A. A. Podgorny

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"



C2

Synthesis of 2-vinylnaphthalene. Yu. S. Zel'kind and
I. A. Arbatova. *Plasticheskie Massy, Sbornik Statei No.*
3, 249-55 (IKRU); *Khim. Referat. Zhur.*, 1960, No. 3, 90.—
2-Vinylnaphthalene (**I**) is synthesized by dehydrogenation
of 2-ethylnaphthalene (**II**). **II** is produced according to the
modified method of Marchetti, but in insignificant amounts.
Better results were obtained by boiling a mixt. of naphthalene,
polyethylated benzene and AlCl_3 . This produced
55% of **I** (calcd. from the amt. of naphthalene used) which
was identified by transforming it into the pterate. **II**
was dehydrogenated by passing the vapors of **II** over cata-
lysts consisting of a mixt. of Cr_2O_3 and MgO or of Al_2O_3
and ZnO . The reaction was carried out at a reduced pres-
sure at 600-700° in an atm. of H_2 . The yield of **I** was
approx. 40-2% of the wt. of **II** and approx. 53-5% of the
wt. of the reaction product. Pure **I** was obtained by distn.
Owing to slight polymerization only approx. 60% of the
crude product was obtained. After recrystg. several times

pure **I** was obtained and identified by transforming it into
the corresponding dibromide. By polymerizing **I** in the
presence of Bz_2O a solid, transparent, slightly yellow poly-
mer was obtained. Polymerization of the impure product
gave the same results. References and patents for produc-
ing **II** and **I** and for polymerizing **I** are given.

W. R. Henn

ASH-14A METALLURGICAL LITERATURE CLASSIFICATION											
SEARCH SYMBOLS											
141080 94	1	2	3	4	5	6	7	8	9	10	11
14 10 80 NO 94	D	N	C	P	R	K	M	H	E	G	T
	0	1	2	3	4	5	6	7	8	9	0

CH

A method for the analysis of terpenes and the composition of Kusnetzovka turpentine. Yu. S. Zal'kind and A. G. Bulavkin. *J. Gen. Chem. (U. S. S. R.)* 9, 309-78 (1939).—The method of Darmois (*C. A.*, 32, 2888) is discussed and shown to be valid only for mixtures of optically active terpenes. Contrary to the results of earlier investigators, turpentine obtained from *Pinus sylvestris* grown in central European Russia contains small amounts of *l*- β -pinene (I). Two com. grades of turpentine, after repeated fractional dist., have an av. compn. of 64% *d*- α -pinene (II), 6% I, 14% *d*- α -carenne (III) and 12% higher-boiling products. II, b_4 54°, d_4^{20} 0.859, n_D^2 1.4000, α/α_0 1.14, α/α_0 1.96 ($\nu = 546.1$ cm, $\delta = 578$ cm, $f = 486.1$ cm, $\epsilon \sim 656.3$ cm), temp. of soln. in Ac₂O 53.8°, I, b_4 104.0-4.5°, d_4^{20} 0.8732, n_D^2 1.4795, $[\alpha]_D - 19.81^\circ$, α/α_0 1.008, α/α_0 1.430. III, b_4 68°, d_4^{20} 0.865, n_D^2 1.4728, α/α_0 1.16, α/α_0 2.17, temp. of soln. in Ac₂O 39°. The presence of I is shown by oxidation with alk. KMnO₄ to give nopinic acid, m. 124-7°, and by conversion to borneol, m. 198-202.5°. Owing to the anomalous values obtained the dispersion coeffs. α/α_0 and α/α_0 possess little value for characterization of mixtures. contg. I and *d*-rotatory terpenes (II, III).


John Livak

20

ASA-SLA METALLURGICAL LITERATURE CLASSIFICATION

E-Z FILE INDEX

E-Z FILE SUBJECT		E-Z FILE SUBJECT	
140002	140003	140004	140005
S	M	H	N
D	U	I	O
W	E	K	R
G	V	X	S
C	F	Z	T
B	P	L	W
A	D	M	H
M	R	N	S
L	O	O	T
Y	Y	Y	Y



		1ST AND 2ND ORDERS		PROCESSES AND PROPERTIES INDEX			
1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
49	50	51	52	53	54	55	56
57	58	59	60	61	62	63	64
65	66	67	68	69	70	71	72
73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88
89	90	91	92	93	94	95	96
97	98	99	100	101	102	103	104
105	106	107	108	109	110	111	112
113	114	115	116	117	118	119	120
121	122	123	124	125	126	127	128
129	130	131	132	133	134	135	136
137	138	139	140	141	142	143	144
145	146	147	148	149	150	151	152
153	154	155	156	157	158	159	160
161	162	163	164	165	166	167	168
169	170	171	172	173	174	175	176
177	178	179	180	181	182	183	184
185	186	187	188	189	190	191	192
193	194	195	196	197	198	199	200
201	202	203	204	205	206	207	208
209	210	211	212	213	214	215	216
217	218	219	220	221	222	223	224
225	226	227	228	229	230	231	232
233	234	235	236	237	238	239	240
241	242	243	244	245	246	247	248
249	250	251	252	253	254	255	256
257	258	259	260	261	262	263	264
265	266	267	268	269	270	271	272
273	274	275	276	277	278	279	280
281	282	283	284	285	286	287	288
289	290	291	292	293	294	295	296
297	298	299	300	301	302	303	304
305	306	307	308	309	310	311	312
313	314	315	316	317	318	319	320
321	322	323	324	325	326	327	328
329	330	331	332	333	334	335	336
337	338	339	340	341	342	343	344
345	346	347	348	349	350	351	352
353	354	355	356	357	358	359	360
361	362	363	364	365	366	367	368
369	370	371	372	373	374	375	376
377	378	379	380	381	382	383	384
385	386	387	388	389	390	391	392
393	394	395	396	397	398	399	400
401	402	403	404	405	406	407	408
409	410	411	412	413	414	415	416
417	418	419	420	421	422	423	424
425	426	427	428	429	430	431	432
433	434	435	436	437	438	439	440
441	442	443	444	445	446	447	448
449	450	451	452	453	454	455	456
457	458	459	460	461	462	463	464
465	466	467	468	469	470	471	472
473	474	475	476	477	478	479	480
481	482	483	484	485	486	487	488
489	490	491	492	493	494	495	496
497	498	499	500	501	502	503	504
505	506	507	508	509	510	511	512
513	514	515	516	517	518	519	520
521	522	523	524	525	526	527	528
529	530	531	532	533	534	535	536
537	538	539	540	541	542	543	544
545	546	547	548	549	550	551	552
553	554	555	556	557	558	559	560
561	562	563	564	565	566	567	568
569	570	571	572	573	574	575	576
577	578	579	580	581	582	583	584
585	586	587	588	589	590	591	592
593	594	595	596	597	598	599	600
601	602	603	604	605	606	607	608
609	610	611	612	613	614	615	616
617	618	619	620	621	622	623	624
625	626	627	628	629	630	631	632
633	634	635	636	637	638	639	640
641	642	643	644	645	646	647	648
649	650	651	652	653	654	655	656
657	658	659	660	661	662	663	664
665	666	667	668	669	670	671	672
673	674	675	676	677	678	679	680
681	682	683	684	685	686	687	688
689	690	691	692	693	694	695	696
697	698	699	700	701	702	703	704
705	706	707	708	709	710	711	712
713	714	715	716	717	718	719	720
721	722	723	724	725	726	727	728
729	730	731	732	733	734	735	736
737	738	739	740	741	742	743	744
745	746	747	748	749	750	751	752
753	754	755	756	757	758	759	760
761	762	763	764	765	766	767	768
769	770	771	772	773	774	775	776
777	778	779	780	781	782	783	784
785	786	787	788	789	790	791	792
793	794	795	796	797	798	799	800
801	802	803	804	805	806	807	808
809	810	811	812	813	814	815	816
817	818	819	820	821	822	823	824
825	826	827	828	829	830	831	832
833	834	835	836	837	838	839	840
841	842	843	844	845	846	847	848
849	850	851	852	853	854	855	856
857	858	859	860	861	862	863	864
865	866	867	868	869	870	871	872
873	874	875	876	877	878	879	880
881	882	883	884	885	886	887	888
889	890	891	892	893	894	895	896
897	898	899	900	901	902	903	904
905	906	907	908	909	910	911	912
913	914	915	916	917	918	919	920
921	922	923	924	925	926	927	928
929	930	931	932	933	934	935	936
937	938	939	940	941	942	943	944
945	946	947	948	949	950	951	952
953	954	955	956	957	958	959	960
961	962	963	964	965	966	967	968
969	970	971	972	973	974	975	976
977	978	979	980	981	982	983	984
985	986	987	988	989	990	991	992
993	994	995	996	997	998	999	1000

The synthesis of an alcohol with two conjugated triple bonds. Yu. S. Zal'kind and M. A. Afitskovich. *J. Gen. Chem. (U.S.S.R.)* 9: 1011-1030. (1939).—On heating 10 g. 2,7-dimethyl-3,5-octadiyne-2,7-diol with 20 g. $\text{Ca}(\text{OH})_2$ at 75°-130°, it is cleaved, giving a mixt. of $\text{Me}_2\text{C}=\text{CH}-\text{CH}_2-\text{CH}=\text{CH}_2$ and traces of 3-methyl-1,7-heptadien-3-ol, $\text{Me}_2\text{C}(\text{OH})\text{C}(\text{CH}_3)\text{CH}_2\text{CH}=\text{CH}_2$ (I). The same results were obtained by using K_2CO_3 or $\text{Ba}(\text{OH})_2$. However, on heating the above reactants for only 30 min., i. e., terminating the reaction before its completion at 120-130°, 17% (theory) of I was obtained. In this case, 46.5% of glycol was recovered. I, b.p. 50-61°, b.v. 72-4°, d₄²⁰ 0.9395, n_D²⁰ 1.4850; MR_D = 33.00. Since I is not very stable, it was exdt. from the reaction mixt., after distg. off most of the $\text{H}_2\text{C}\equiv\text{CH}_2$ and Me_2CO , by NH_3 , Ag_2O , and the resulting ppt. was decompd. (cold) with 20% H_2SO_4 under an Et_2O layer (several days required). The I formed immediately dissolved in the Et_2O layer, and was recovered from it in the usual manner. $\text{H}_2\text{C}\equiv\text{CH}_2$ glycals with 2 conjugated triple bonds on heating with K_2CO_3 , $\text{Ca}(\text{OH})_2$, or $\text{Ba}(\text{OH})_2$, decomp., forming 2 mol. of ketone and $\text{H}_2\text{C}\equiv\text{CH}_2$ or 1 mol. of ketone and $\text{H}_2\text{C}\equiv\text{CH}_2$ alc.

A. A. Podgorny

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

Synthesis of glycols of the bicyacetylene series. Yu. S. Zal'kind and I. M. Overdijkell, *J. Gen. Chem. (USSR)*, **4**, 971-4 (1939); cf. *C. A.*, **31**, 4289. The previously described method for the synthesis of bicyacetylene glycols was used in the presn. of Ni(*t*-hydroxycyclohexyl)-acetylene (II_1 , m. 131.2-4.2°, from *t*-hydroxycyclohexyl-acetylene). The yield was almost quant. I_1 hydrogenated in the presence of Pt black, added 8 H atoms, forming *t*,*t*-di(*t*-hydroxycyclohexyl)butane, m. 91.8 (22.8%). In the presence of colloidal Pd, the hydrogenation after addn. of 8 H atoms proceeded very slowly. Stopping the reaction at this stage yielded, probably, *t*,*t*-di(*t*-hydroxycyclohexyl)-*t*,*t*-butadiene, m. 101-4°. The application of this reaction to the synthesis of bicyacetylene derivs. of OH compds., using secondary acetylene ales., was also investigated. For this purpose, the following ales. were

prep'd. by the Teterov and Lymanova method (cf. *C. A.*, **44**, 8003) from the corresponding aldehydes: t -MgC₂-C₆H₅MgBr and ales.: *t*-butyne-*1*,*1*, m. 107-0°; *t*-pentyn-*1*,*1*, m. 124.4°, d₂₀²⁰ 0.8851, n_D²⁰ 1.43447, MRn 24.74; *t*-hexyn-*1*,*1*, m. 140-1°, d₂₀²⁰ 0.8704, n_D²⁰ 1.43297, MRn 20.38; and *t*-heptynyl-*1*,*1*,*1*,*1*,*1*,*1*,*1*, PhCH(OH)C₂CH₂, m. 114-17°, d₂₀²⁰ 1.0215, n_D²⁰ 1.5201, MRn 39.815. In all cases the yield was 20-35% (theory). The condensation of these ales. yielded, in very small amounts, 3,3-*cyclododec-2,7-diol*, in 2 isomeric forms, m. 107.0° and 107.0°; 4,9-*deoxydodec-4,9-diol*, [$\text{MeCH}_2\text{H}_2\text{OHC}(\text{CH}_2)_8\text{CO}_2\text{Me}$], m. 164.6°; 3,7-*deoxydodec-4,9-diol*, [$\text{Me}_2\text{CH}_2\text{H}_2\text{OHC}(\text{CH}_2)_8\text{CO}_2\text{Me}$], m. 159.0°; and 1,6-*diphenyl-2,6-heptadieno-1,6-diol*, [PhCH(OH)C₂CH₂], m. 132-3°. The condensation probably proceeded through the formation of free radicals by removal of a H atom with subsequent condensation of free radicals. The formation of free radicals in the case of secondary ales. is retarded because the pos. inductive effect of Me group, which promotes this process, is balanced by the neg. effect of OH, whereas in the case of tertiary ales. the effect of 2 Me groups is only partially balanced by 1 OH group. This explains the low yields of bicyacetylene derivs. of OH compds. A. A. Podgorny

C4

The reaction between acetylenedimagnesium dibromide and carbonyl compounds. Yu. S. Zel'kind and S. M. Labazov. *J. Russ. Chem. Phys.* S. R. 37, 1527-32 (1969). — The influence of the structure of CO compds. on the condensation of aldehydes and ketones with $(\text{AcMgBr})_2$ (I) in the synthesis of acetylene γ -glycols by the lotrich method is studied. To this end, I, prep., from 2 g. Mg, 12 g. BrC_2 and 10 ml. ether, is treated at 0° with a definite quantity of an aldehyde and ketone for measured periods of time and the reaction mixt. is decomposed with H_2O (contg. H_2S). The vol. of liberated C_2H_2 gives the amt. of unaltered I and indirectly the C_2H_2 used in the reaction. The exact content of I in the reagent was detd. by expelling the ether and decomposing the residue with H_2O . It was 91.5 ± 10 ml. C_2H_2 , corresponding to 98.8% I. This figure was used in calcg. the results of the condensation reactions. Furthermore, since some C_2H_2 is dissolved by ether in the I reagent, its mean content was detd. by decomposing the reagent with water without the preliminary removal of the ether and heating the mixt. on a water bath. A vol. of 1240 ml. C_2H_2 was collected of 331 ml. in excess of the amt. liberated in the decomp. of I reagent with the previous expulsion of the ether. This C_2H_2 amt. was deducted from the C_2H_2 formed in the decomp. of the condensation reac-

tion mixt. In the condensation with MeCO , MeCOEt , MeCOPr , BrCO , MeCOPh and PhCO the reaction proceeds rapidly in the first 15 min. with the decomp. of 37.8% I and reaches a max. of 57.8% in 1 hr. This limit is not exceeded by longer duration of the reaction (1 hrs.) or by the use of a large excess of ketones (10 times). In the presence of I excess the condensation proceeds nearly quantitatively. The results show that the reaction is not reversible, hence the incomplete condensation is not caused by a reaction equil. in conformity with the law of mass action. It is probable that the reaction stops because of the occlusion of I by the insol. Mg complex formed in the reaction. The reaction proceeds in 2 stages: $\text{MeCO} + \text{I} \rightarrow \text{MeC}(\text{OMgBr})\text{C}\equiv\text{CMgBr}$, $\text{MeC}(\text{OMgBr})\text{C}\equiv\text{CMgBr} + \text{MeCO} \rightarrow (\text{MeC}(\text{OMgBr})\text{C}\equiv\text{C})_2$. This is confirmed by the formation of a glycol tetramethyl-2-butene-1,4-diol and alc. ($\text{MeC}(\text{OH})(\text{CH}_2)_3\text{CH}_2\text{OH}$). In general, the velocity of condensation decreases with increasing mol. wt. of ketones. Aldehydes (AcH , BzCHO , PrCHO and BzH) react with I more rapidly to a max. of 70-100% decompr. (AcH).

CAT

preparation of substituted diphenylacetylenes. Yu. S. Zal'kikh and B. M. Fudylev. *J. Gen. Chem. (U.S.S.R.)* 9, 1725-81 (1939).—In a previous paper the authors described the synthesis of PhC₂CCl₂CPH from PhC₂Cl₂ and NH₂CuCl. Similarly tertiary alcs. with acetylenic radicals form corresponding diacetylenic glycols, a reaction which is unsuccessful with secondary acetylenic alcs. (low yields). ρ -Methyl-, ρ -chloro-, ρ -bromo-, and ρ -nitrophenylacetylenes were subjected to this reaction to check on the electronic theory of Ingold and Robinson. The following yields and time of complex formation were noticed: ρ -tolyl 0%, immediate; Ph 75%, 25-30 min.; ρ -C₆H₄Cl₂, 50%, 30-5 min.; ρ -BrC₆H₄, 45%, 45-80 min.; ρ -NO₂C₆H₄, 25%, 90-120 min. *tert*-BuC₂CH₂ yielded di-*tert*-butylacetylene. ρ -Tolylacetylene was prep'd. according to Kunckell and Goetsch (*Ber.* 43, 2860 (1900)) by the action of Na on ρ -MeC₆H₄CCl₂CHCl, obtained from MeC₆H₄COCl₂ and PCl₅. ρ -Tolylacetylene (3 g.) was added dropwise to a soln. of 10 g. of CuCl, 16 g. of NH₃, and 0.15 cc. of concn. HCl in 48 cc. of H₂O at 0°-65°. The formation of a solid complex was immediate. Heating lasted for 2 hrs. A current of alc. was continuously passed through the mist. When treated with ether the complex decompd. A 90% yield of di- ρ -tolylacetylene, m. 183-3.2°, was obtained. A mixt. of 85 g. ClC₆H₄CCl₂AlCl, bp.-110-118°, and 8 g. ClC₆H₄CCl₂Cl₂, bp.-110°, was prep'd. from 94 g. of ρ -chlorosobophenone and PCl₅. These chlorides were boiled 3 hrs. with 800 g. of 25% KOH in alc. After allowing the mixt. to stand for 24 hrs. the product was treated with H₂O₂, extd. with ether, dried and vacuum-distd., bp.-10 85-95°, m. 41-3°. Di- ρ -chlorophenylacetylene was obtained in 80% yield, m. 238°. Di- ρ -bromophenylacetylene, m. 203-4°, was prep'd. similarly to the Cl deriv. ρ -Nitrophenylacetylene was prep'd. according to Drewsen, *Ann.* 312, 164. It was condensd. as in previous expts., yielding di- ρ -nitrophenylacetylene, m. 249-5°.

D. Acetyly


Preparation of cyclohexanol from cyclohexene. Yu. N. Zel'kind and I. V. Markov. *J. Applied Chem. (U. S. S. R.)* 12, 437-9 (in French, 429) (1939).—Mixing 1 mol. of cyclohexene with 1-1.5 mols. of 70-85% HgSO₄ at 0°-6° for 30 min., subsequent diln. of the reaction mixt. with 5 vols. of water, steam distg., settling out with NaCl and drying with K₂CO₃ yielded, after fractionation, 60-64% of cyclohexanol. The cyclohexene obtained from petroleum yielded less cyclohexanol, probably because of impurities. A Ag₂O catalyst slightly increased the yield and decreased the tar formation. A K₂Fe(CN)₆.3H₂O catalyst had no effect.


Investigation of the Malter effect. M. S. Kostyan, Ya. E. Zal'kind, and M. E. Zal'kind. *Izv. Akad. Nauk SSSR, Ser. Fiz.*, No. 10, 1940, 45-55 (in Russian); cf. Malter, *J. Phys. Rev.* **50**, 48, 1936; — (11).

The abnormally high secondary electron emission occurring from Al_2O_3 and the persistence of the after-effect observed at the end of the primary bombardment were further confirmed by forming on 4 different dielectrics Al_2O_3 , mica, Ca and NaF , thicks, and MgO all on a grounded Ni support. The basic condition for the after-effect appears to be suppression of the electron surface current, with well-sustained mica, the Malter current intensity drops only by 2-3% in 3 min. On the other hand, in each case with insulators possessing a low dielectric loss factor, with insulators possessing a high dielectric loss factor, because of ionization of the dielectric, especially with powders (Al_2O_3), because of ionization of the surface, sudden changes in occur at least part of the extent of 10^{-5} in either direction; in this case, a maximum current for up to 4-5 hrs, with only slight fluctuations and then suddenly disappears. (2) At a given primary current intensity, the Malter current increases steadily with the collector voltage V_C ; this holds also for the after-effect ($I = I_0$), which indicates that at least part of the electrons are emitted by the dielectric. There are evidently 2 processes present: emission from the dielectric spontaneously, that is without primary bombardment, without replacement by electrons from the metal surface, and port, resulting in raising the potential of the surface, and emission with attachment, the second process being stationary state. Indication for the first process is seen in the observation that the Malter current "revives" when the collector potential is again applied after having been lowered to $V_C = 0$, discrete centers in which the potential is raised, assumed to be present on the dielectric, evidently persist at $V_C = 0$ and on its reapplication give rise to emission by the first mechanism. This is confirmed by the first measurements of the potential reach of the primary beam, which is determined by the center of repulsion of the electron current, that is, by the potential barrier at $V_C = 0$ separating the center of repulsion subsequently "secondary" revival. (3) Flowing on glassed films, with or without electrodes, on Al_2O_3 powder, with or without electrodes, with $I = I_0$ the currents with nearly the same value of V_C as in the case of the dielectric, and, tends to decrease with rising I . In this case, the current density distribution curves at high current densities, $|J| \sim V_C^{-1}$, show most of the electrons to have left the cathode to a negligible fraction having velocities corresponding to a potential drop across the dielectric layer. Thus in the highest depths of the film, the alternative interlayer current, namely that electrons are emitted from the metal support but lose energy on traversing the film, is contradicted by the curve of angular distribution showing the majority of electrons to be emitted perpendicular to the surface in absence of any noticeable scattering such as would result from loss of energy through collisions. At low I , there is a substantial number of electrons having full velocity; this indicates that in this case most electrons are emitted from the metal support. N. Thor.

Thom

APPROVED FOR RELEASE: 09/19/2001 CIA-RDP86-00513R001963710009-5"





10

The addition of hydrogen to acetylene compounds. XXXII. Catalytic hydrogenation of an alcohol with a double and a triple bond. Yu. S. Zal'kind and N. D. Khuckova. *J. Gen. Chem. (U. S. S. R.)* 10, 621-6 (1940); cf. *C. A.* 34, 3871.— $\text{CH}_2\text{CH}(\text{CH}_3)\text{Br}$ and Mg with MeLiCO give 77.5-88% $\text{MeEtC(OH)C(=O)CH}_2\text{CH}_3$, bp 75-8°, bp 81-2°. When this is hydrogenated over Pt black it easily adds 4 H atoms, then in about double the time it adds 2 more. A very slow reaction with H follows, probably due to reduction of the OH group. When colloidal Pd deposited on starch is used, the addn. of the last 2 H atoms takes 3-4 times as long as addn. of the first 4, and there is no further reaction. Addn. of the first 2 H atoms goes almost entirely on the triple bond, giving a compd. which reacts with maleic anhydride to form crystals m. 136-7°. The next 2 H atoms add to any of the available double bonds, giving a mixt. of unsatd. alcs. When 6 atoms of H have been added, the product is MeEtBuCOH . H. M. Leicester


COMPOUND ELEMENTS

MATERIALS INDEX


ABRILIA METALLURGICAL LITERATURE CLASSIFICATION

FROMI SYNTESIZA

VOLUME NO.	ISSN NO.	PUBLICATION DATE	EDITION	ECONOMIC INDUSTRIAL												
				1	2	3	4	5	6	7	8	9	10	11	12	13
14	15	MAY 1963	1	0	0	0	0	0	0	0	0	0	0	0	0	0



Dehydration of alcohols containing two triple bonds and the products of this reaction. Yu. S. Zal'kind and E. F. Zelberg. *J. Gen. Chem. (U. S.S.R.)* 10, 1739-1812 (1940). - Prepn. of hydrocarbons of type R-C=C(CR)- was attempted by dehydration of the corresponding diethynyl alcohols, $(\text{PhC}_2\text{C})_2\text{C}(\text{OH})\text{Me}$ (I), from PhC_2CMgBr and EtO_2Ac , was dehydrated by H_2SO_4 , H_3PO_4 , KHSO_4 , $(\text{C}_6\text{H}_5)_2\text{N}$, ρ -Mc₆C₆SO₃H, yielding in all cases only the polymer of the desired hydrocarbon; *unsym-bis(phenyl-ethynyl)ethylene* (II). All work was done in an inert atm. due to sensitivity of the products. The latter were brownish powders which were fractionated by benzene-Et₂O prts., into a series from the dimer to the octamer of II. The dimer and trimer were unsat'd. and tended to polymerize further on standing, while the higher polymers were stable. The following formulas are proposed for the lower and higher polymers:



Hexyldis(phenylethynyl)carbinol (**III**). From EtMgBr, PhC≡CH and Et heptanoate, was prep'd. in order to study the effect of a larger group R. However, dehydration of **III** by 60% H₂O₂ also yielded polymeric material only, as an alc.-insol. brown solid, apparently a tetramer of the tested hydrocarbon. **III** m.p. 65-5.5° (from petr. ether). G. M. Kosolapoff

G. M. Kowalewski

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

The addition of hydrogen to acetylene compounds. XXXIII. Synthesis and catalytic hydrogenation of hydroxy acids of the acetylene series. Yu. S. Zal'kind and B. I. Mikhant'ev. *J. Gen. Chem. (U. S. S. R.)* 11, 92-8 (1941); cf. *C. A.* 34, 7847. When CO_2 is passed through Me_2CO and the mixed vapors are passed into $\text{BrMgCl}\text{C:CMgBr}$ (I) at -8° , 0.75% 1,1-dimethyl-2-butyn-1-ol-4-carboxylic acid (II), decomps., on vacuum distn., is obtained. When $\text{Me}_2\text{CO}(\text{OMgBr})\text{C:CMgBr}$ (III) is treated with CO_2 at 8° , the yield of II is 1.25%. This can be raised somewhat if the CO_2 is passed into the mixt. for 5 days. Treatment of III with ClCO_2Me gives 1.5% II. The Ag, Cu, Ba and Ca salts of II are insol. The Ag salt is explosive. The amide m. 72-3°. In similar reactions I, Ph_2CO and CO_2 give 0.7% 1,1-diphenyl-2-butyn-1-ol-4-carboxylic acid (IV), m. 78-80°, decomps., 127° $\text{PhC}(\text{OMgBr})\text{C:CMgBr}$ (V) and CO_2 give 3.6% IV, and V with ClCO_2Me gives 0.44% IV. The Ag, Cu, Ca, Hg, Si and Pb salts of IV are insol.; the amide m. 96-7°. Hydrogenation of II, IV, and their amides over Pd and Pt adds 4 H atoms. The rate of addn. over Pt is regular, but over Pd the last 2 H atoms add more slowly. This effect is not as marked as in the acetylene γ -glycols however.

H. M. Lester

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"

Cd


Addition of hydrogen to acetylenic derivatives. *III* (2). Catalytic hydrogenation of phenylated butyne-diol in the presence of colloidal palladium. *VII*, *S.* Zal'kind and A. P. Ivanov. *J. Gen. Chem. (U. S. S. R.)* 11, 885 (1941); *cf. C. A.* 35, 5529. — The authors made a comparative study of the hydrogenation of 10 acetylenic glycols in the presence of colloidal Pd under controlled conditions. $\text{BrMgC}_2\text{CMgBr}$ (from 30 g. Mg) was treated, with cooling and stirring, with 60 g. McCOPh and 15 g. abs. EtOH in dry Et_2O , let stand 24 hrs., treated with moist Et_2O , the aq. layer treated with weak H_2SO_4 , and Et_2O ; both Et_2O exts. were dried and the Et_2O distd.; the 1st ext. yielded a mixt. of *trans*-dimethylphenylbutynediol and *2-phenyl-3-butyn-2-ol* (*I*); the latter was dissolved in warm petr. ether, the solvent distd., and the residue steam-distd., the distillate satd. with K_2CO_3 and extd. with Et_2O . The 2nd Et_2O ext. treated as above also yielded *I*, giving a total of 21.5 g. *I* (20.4%), m. p. 48.5–50° (from petr. ether). EtMgBr (from 14.4 g. Mg) in ether was treated with 24 g. dimethylethynecarbinol in Et_2O with cooling and stirring, followed by 21 g. EtOH in ether; after 2 days the mixt. was treated with moist Et_2O and ice water, and the aq. layer treated with dil. H_2SO_4 and extd. with Et_2O . On distn. of the Et_2O both exts. gave crystals of *4-methyl-1-phenyl-2-butyne-1,4-diol*, (*II*), which was recrystd. from benzene and Et_2O (20 g. Et_2O), m. p. 73.6–74°. EtMgBr (from 3.0 g. Mg), treated with 0.01 g. phenylethynecarbinol with cooling and stirring, warmed for 0.5 hr. on a water bath, then treated with 0.1 g. PhCO in ether and let stand for 3 days, heated for 0.5 hr. on a steam bath and worked up as

above, gave 11 g. (67.8%) *1,1,4-triphenyl-2-butyne-1,4-diol* (*III*). *III* was also prep'd. from EtMgBr (from 2.5 g. Mg), 10.1 g. diphenylethynecarbinol and 5 g. EtOH in 64.3% yield. *III*, m. p. 110–111°. *2-Methyl-3-phenyl-3-hexyn-2-ol* (*IV*), prep'd. as described above from 14.4 g. Mg, 60 g. EtOH , 24 g. dimethylethynecarbinol and 27 g. McCOPh , in 70.2% yield (35 g.), m. p. 81.5–82°. *1,1,4-Triphenyl-2-butyne-1,4-diol* (*V*), prep'd. as described above from 4.8 g. Mg, 22 g. EtOH , 14.6 g. I, and 15 g. PhCO in 34.6% yield (9.4 g.), m. p. 125.5–126°. *3-Methyl-1-(9-hydroxy-9-fluorenyl)-1-butyne-2-ol* (*VI*), prep'd. from 37 g. fluorene, 18 g. Mg, 98 g. EtOH and 34 g. dimethylethynecarbinol as described above, was obtained in 62.7% yield (34 g.), m. p. 111.5° (from EtOH). The 5 glycols and 5 previously known ones: diphenyl-, tetra-

methyl-, *trans*-dimethyldiphenyl-, *syn*-dimethyldiphenyl-, and tetraphenylbutynediol, were hydrogenated alike, by using 0.01 g. mol. of the glycol in 50 cc. Et_2O and 2 mg. colloidal Pd. All but *V* and tetraphenylbutynediol were run at 10° and 743 mm.; the remaining 2 were run at 17.5° and 743 mm., and 20° and 734.5 mm., resp. In addition, exts. were also run with 4 mg. Pd catalyst, and all new glycols (*II*–*VI*) were also run with 6 mg. Pd. The reaction consts. ($\times 10^3$) for 2 mg. Pd were found to be: dihexynediol 0.5, *II* 0.16, *III* 3.3, tetramethyl-

butynediol 0.6, IV 3.6, *syn,syn*-dimethylidiphenylbutynediol 1.37, V 0.80, diol 3.4, *syn*-dimethylidiphenylbutynediol 0.43, VI 4.09. For 4 mg. Pd the tetraphenylbutynediol 0.43, VI 4.09. For 4 mg. Pd the values were 13.2, 11.2, 7.0, 16.1, 10.4, 8.5, —, 2.1, 0.6, and 11.3, resp. Hydrogenation with 6 mg. Pd was run mainly to observe any rapid change of rate of hydrogenation after addition of the 1st 2 H atoms; in all cases a definite decrease of hydrogenation rate was observed after formation of the ethylene glycol-type structure. Although the reaction, in general, is not strictly a 1st-order reaction, and K varies in the course of its progress, the av. values of K (given above) were used, because they are convenient measuring comparison standards. The products of hydrogenation of II-VI were isolated as follows: the Pd was filtered off, the RtOAc distd. in vacuo, the residue dissolved in Et_2O , dried, the Et_2O distilled off and the residue distilled, if possible. II yielded 4-methyl-1-phenyl-2-pentene-1,4-diol, b.p. 154°; III yielded 1,1,4-triphenyl-2-butene-1,4-diol, a viscous liquid which could not be distd. without decompr.; IV yielded 2-methyl-3-phenyl-3-hexene-2,5-diol, b.p. 175-8°; V yielded 1,1,4-triphenyl-2-pentene-1,4-diol, m.p. 75-6° (from benzene); VI gave 3-methyl-1-(*n*-hydroxy-9-fluorend)-4-butene-3-ol, m.p. 182-3° (from EtOH). In general, disubstituted and secondary-tertiary glycols hydrogenate more readily than do the tertiary glycols of comparable mol. wt. Substitution of Ph groups for Me groups leads to a slower hydrogenation rate. Closure of 3 Ph groups into a fluorene nucleus causes a great increase in hydrogenation rate, showing the effect of steric structure of the mol.

(3) M. Komlapati



C. G. LEWIS

Addition of H to acetylene derivatives in the presence of Pd or Pt. **LXXXVII.** Catalytic hydrogenation of acetic esters of acetylenic esterified. Yu. S. Zel'kind and A. Kachinskii. *J. Gen. Chem. (U. S. S. R.)* 12, 649-62 (1942) (English summary); cf. *C. A. 36*, 37109— $\text{Br}_3\text{MgC}(\text{CMe})_2$ (from 24 g. Mg) was treated with 80 g. benzoquinone in 200 cc. dry Et_2O , let stand 3 days, heated to 60° for 3 hrs. and treated with dil. H_2SO_4 to yield 2,3,5-triethoxy-3-acryloyl-1,3,5-trisubstituted (64%), m. 218-219° (from Et_2OH); tetrasuccinate (by heating with AgO and NaOAc in the presence of pyridine), m. 185.5-187° (from MePh). The latter hydrogenated in the presence of colloidal Pd at 17° in Et_2OAc gave the corresponding ethylenic compound, $\text{C}_6\text{H}_{10}\text{O}_4$, m. 181.5-183° (from MePh). After this addition of 2 H the reaction slows down greatly with Pd. However, hydrogenation over Pt black in Et_2OAc yielded 1,3,5-triethoxy-1,3,5-trisubstituted tetrasuccinate, m. 187-188° (from MePh). G. M. K.

11

ASA-SEA METALLURGICAL LITERATURE CLASSIFICATION

卷之三

APPROVED FOR RELEASE: 09/19/2001


CIA-RDP86-00513R001963710009-5"

Plasticizers from naphthenic acid esters. Vn. S. Zalikidze,
V. P. Gognadze, and N. K. Iremadze (Tbilisi Chem.
Inst.), *Bull. Acad. Sci. Georgian S.S.R.*, 4, 187-92
(1943) (in Georgian and Russian). - The acids were extd.
from two kerogen (20% 100° and 150-160°) and two oil
(65-100° and 150-210°) fractions of Taku petroleum
distd. under 3.0 mm., and esterified for 21 hrs. on a water
bath with 1% p-toluenesulfonic acid as catalyst with
EtOH (4-fold excess) and with glycerol (2-fold excess),
in the latter case dild. with gasoline which is subsequently
distd. off. From the kerosene acids one obtains Et esters
m. -72, -73; glycerol esters m. -75, -21; from the
oil acids, Et esters m. -60, -48; glycerol esters m. -50,
-52°. By volatility tests (loss of wt. in air after 10 days),
Et esters of the lower fractions are too volatile for use as solvents,
Et esters of the higher fractions, particularly glycerides, have
the right low volatility. No turbidity or pptn. occurs on
mixing with cellulose ester solns. A cellulose lacquer
prep'd. with naphthenic acid glyceride as plasticizer proved
satisfactory. N. Thom

20

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"



Azo derivatives of the phenanthrene series. Yu. S. Zal'kind and Z. I. Khefser's (*Leningrad Chem. Tech. Inst. J. Gen. Chem. (U.S.S.R.)*, **15**, 306-74 (1945) (English summary)—Eight azo dyes based on phenanthrene were prepared, and their spectra are presented. The 3-phenanthryl nucleus causes but slight deepening of color as compared with C_6H_5 , while the 2-phenanthryl radical causes substantial color deepening. Phenanthrene (100 g.) was treated at 143° over 20 min. with 51 g. concd. H_2SO_4 , after which the mixt. was heated for 20 min. longer; after pouring into 1 l. 20% NaCl and neutralization with NaOH the mixt. was allowed to stand for 24 hrs., when the ppt. was filtered off, recrystallized from 1.5 l. water, dissolved in 2 l. H_2O , and, after addition of a little HCl, was treated with a soln. of 18 g. BaCl₂. The pptd. Ba salt was thoroughly washed with hot water and the residue collected as Ba-2-phenanthrenesulfonate (15.5%). The filtrates on concn. and treatment with 5 g. BaCl₂ gave 10-11 g. more of the above Ba salt contaminated with some of the 3-isomer; repetition of the procedure yields an unspecified amt. of the pure 2-isomer. All filtrates were treated with H_2SO_4 , neutralized, after filtration, with KOH and treated with KCl while hot to yield 32 g. of the K-3-phenanthrenesulfonate; addnl. aints. of this acid were obtained as the Ba salt from the filtrate by addn. of BaCl₂; the total yield of the 3-isomer was 31.0%. Fusion with an equal mixt. of KOH-NaOH at 250-300° gave 87% 3-phenanthro (from the 3-sulfonic acid, as K salt), m. p. 120-1°. Fusion of Ba-2-phenanthrenesulfonate with 3 parts of KOH at 310-320° gave 81.5% 2-phenanthro (from lignin). Coupling of these phenols with the appropriate diazonium soln. in the presence of NaOH gave the following dyes: *4-phenyl-2-phenanthro*, m. p. 145-6°; following dyes: *4-phenyl-2-phenanthro*, m. p. 178-80° (from EtOH), deep red (89.1%); *1-phenyl-2-phenanthro* (from EtOH), deep red (no yield given); *4-(2-naphthylazo)-2-phenanthro*, red crystals (no given); *4-(1-naphthylazo)-2-phenanthro*, m. p. 193.4°%; *4-(1-naphthylazo)-2-phenanthro*, m. p. 180-82° (from EtOH), red crystals (88.1%); *1-(1-naphthylazo)-2-phenanthro*, m. p. 229-9° (90.8%); *1-(1-naphthylazo)-2-phenanthro*, gives a yellow color in dil. H_2O ; *3-phenanthro* (2 g.), solns., no m.p. given (91.9%); *3-phenanthro* (2 g.), solns., no m.p. given (73.4%). *2-phenanthrylamine*, m. p. 85°, HCl salt, no m.p. given. Diazotization of *2-phenanthrylamine* and treatment with 2-naphthol in NaOH gave *1-(3-phenanthrylazo)-2-naphthol*, red, no m.p. given; coupling with 3-phenanthro (as above) gave *4-(3-phenanthrylazo)-2-phenanthro*, red powder, does not melt at 300° (from naphthine).

1980-81 EDITION OF THE MEDICAL LITERATURE CLASSIFICATION

ASH-SEA METACOMMERCIAL EXCHANGE

CIA-RDP86-00513R001963710009-5"

151 325 150 44595
PROCESSED AND SERIALIZED MARCH 1967
100 AND 6TH COPIES

Condensation of bis(phenylbutyndial with phenol, methylene-2-chloro-1-phenylidene, m. 157° (cf. Wieland Yu. S. Zal'kind, V. I. Teterin, and S. G. Kurnetsov and Klues, C.A. 23, 3690), and 3 unidentified products m. (Leningrad Chem. Tech. Inst.), J. Gen. Chem. (U.S.S.R.) 17-8 (4.5 g.), m. 194-5° (0.2 g.), and m. 105° (0.4 g.); (1) 15, 433-98 (1945); English summary). — 1-C(OH)-0.4 g. of the chloride was added to a soln. of 0.1 g. K in 2 g. PhCl (1) (30 g.), 37.5 ml. PhOH, 60 cc. benzene and 0.4 g. molten PbOII and heated slowly to 230° for 1 hr.; after dry H₂NCH₂SO₃H II were refluxed for 2 hrs. with continuous treatment with alk. water, extn. of the ppt. with EtOH, and removal of water by means of a Stark-Dean type collector, cryst. of the residue from benzene-petr. ether there was in which 3.35 cc. H₂O was collected at the end of the reaction; obtained a product identical with III, m. 221°, above. After cooling and diln. with benzene, the cryst. and Boiling of this in MeOH in the presence of sulfamic acid the liquid portions were steam-distd. to remove the solvent. There were obtained 21.7 g. cryst. matter and 10.7 g. red failed to effect any isomerization. IV on boiling with alc. resin. Prolonged recryst. from benzene, CHCl₃, and KOH is transformed into 2 substances which were unidentified. ligroin gave the following products: 21.5 g. 1-diphenylmethylene-2-p-hydroxyphenyl-1-phenylidene, m. 253° colorless, m. 250° (from EtOH), then EtOH-MeCO), and orange, m. 208-19°, with the former transforming into a product m. 208-14°, on heating above the m. p. The condensation of I with PhOH was also conducted in the presence of the following catalysts: activated Chasoviansk clay, H₂SO₄-AcOH, and glacial AcOH. The 1st catalyst gave results similar to sulfamic acid while the 2nd catalyst gave only II, m. 203°, in 98% yield (crude); the 3rd catalyst gave only a resinous, ill-defined mixt. of some transformation products of the glycol without condensation with PhOH. G. M. Kowalewff

8°. III was prep'd. by an alternate method for identification: 10.8 g. I in 150 cc. EtO was treated with 22.5 g. PCl₅ at 1° over 6 hrs., stirred for 2 hrs. at 0° and for 4 hrs. at room temp. to yield, after removal of the solvent, hydrolysis, and crystn. from ligroin 1.0 g. 1-diphenyl-

ASA-SLA METALLURGICAL LITERATURE CLASSIFICATION

150-151 325 150 44595

150-151 325 150 44595

RELATION:


EXCERPT SUMMARY

150-151 325 150 44595

COMMON ELEMENTS		PROCESSES AND PROPERTIES INDEX												COMMON ELEMENTS													
		1ST 100 JUN 1950 191												1ED 100 JUN 1950 191													
CH ₃		Isomerization of an alcohol with two double bonds. Yu. S. Zal'kina and A. I. Kulikov (Leningrad Chem. Tech. Inst.). "J. Russ. Chem. (U.S.S.R.)" 15, 643-9 (1945) (English summary).—The propenyl and hydrogenation of β -methyl(α -methyl)cetobutene (I) were studied. Hydrogenation results apparently in 1-phenyl-2,4-pentadienes-1-ol, which isomerizes into 1-methyl-1,4-pentadiene-3-ol (II), and 1-methyl-1,3-pentadiene-3-ol (III). The starting material was prepd. conventionally from Bril and CH ₃ :CHC:CMgX, with the best (73%) yield resulting from addn. of Bell to the Mg compnd. at -10° and letting the mixt. stand for 4-6 hrs. at -10°, and 12 hrs. at room temp. I, b. 112°, b. 125°, b. 131°, b. 137°, b. 140°, b. 142°, d ₄ ²⁵ 1.0453, n _D ²⁵ 1.57407. Hydrogenation of I in KOAc in the presence of colloidal Pd gave the diene alc. mixt., b. 111-12°, d ₄ ²⁵ 0.9024, n _D ²⁵ 1.54036; it remains quite unchanged on treatment with alc. KOH and its association gave HCHO, BaH, citric acid, and aldehyde. The diene alc. mixt. (8 g.) and 8 g. Cl ₂ CO ₂ H in 8 cc. glacial AcOH, kept in a sealed tube for 22 hrs. at room temp. and for 4 hrs. at 100°, yielded 2 fractions of <i>cis</i> -trichloroacetate, 0.3 g., b. 94-105°, n _D ²⁵ 1.54052, and 0.7 g., b. 101-5°, n _D ²⁵ 1.53211; better results were obtained when distn. was omitted. Hydrolysis of the crude ester, after washing with water, with alc. KOH at reflux yielded 3 fractions of III, 0.7 g., b. 117°, 0.7 g., b. 117-30°, n _D ²⁵ 1.52307, and 0.5 g., b. 120°, formed apparently from iso-	CH ₃																								
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC												SEARCH SYSTEMATIC													
SEARCH SYSTEMATIC	SEARCH SYSTEMATIC	SEARCH SYSTEMATIC																									

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963710009-5"




CH
Action of some inhibitors in catalytic hydrogenation in
the presence of palladium. Yu. S. Zal'khol'd and Kh. V.
Hal'yan. Trudy Leningrad. Tekhnol. Inst. im. Leningrad.
Sovia 1946, No. 12, 73-86. --The inhibiting action of

NH_4CNS , $(\text{NH}_3)_2\text{CS}$, $\text{Pb}(\text{OAc})_2$, and $\text{Cu}(\text{OAc})_2$ during
hydrogenation with colloidal Pd of $(\text{CC}(\text{OH})\text{Me})_2$ (I),
 $\text{Me}_2\text{C}(\text{OH})\text{ClCH}_2$ (II), and $\text{Me}_2\text{C}(\text{OH})\text{CH}_2\text{CH}_3$ (III)
in general decreases in the above order. The Pb salt is
more active with alcs., and the Cu salt with the glycol.
II is less affected by the inhibitors than I, but III some-
what more than II and less than I. Ethylenic alcs. absorb
II fastest when *in situ* *nascendi* from the acetylenic ones.
KCN, Na_2AsO_4 , $\text{Na}_2\text{AsO}_4 \cdot \text{H}_2\text{O}$, CoSO_4 , and NiSO_4 are inactive
as inhibitors. The inhibiting action is believed to be due
to an interaction of the inhibitor with the org. compd.
rather than with the Pd. Exptl. data are tabulated.

Kitty Lus

Condensation of *syn*-tetraphenylbutynediol with β -*tolyl*. Yu. S. Zal'kind, V. K. Teterin, and S. G. Kuznetsov (Leningrad Chem. Tech. Inst.), *J. Gen. Chem. U.S.S.R.* **16**, 611-20 (1940); cf. *C.A.* **40**, 17058. *Tetraphenylbutynediol* (**I**) (9.8 g.), 10.8 g. *p*-*tolyl*, 26 cc. benzene, and 0.1 g. anhyd. sulfamic acid were refluxed 5 hrs., using a reflux trap to collect the resulting water, of which 0.72 cc. was collected; the residual mass was distd. with steam and the residual red oil was dried by azeotropic distn. with benzene. On cooling the benzene soln. there was obtained 2.1 g. *1-diphenylmethylenec-3-(p-toloxo)-3-phenylindene* (**II**) m. 215° (from benzene), while the benzene mother liquor on diln. with ligroin gave a mixt. of 2 substances, sep'd. by leaching with Et₂O, which left behind 2.5 g. *6-methyl-2,2',3'-triphenoxy-1',2'*_{3,4}-indanethrene (**III**), m. 222-4° (from ligroin); repeated crystl. of the Et₂O-sol. material from benzene-ligroin gave 2.3 g. *2,2,5,5-tetraphenyl-3-(p-toloxo)-2,5-dihydrofuran*, m. 136° (from petr. ether); concn. of the mother liquor and prolonged standing gave 0.02 g. *tetraphenylbatiene*, m. 226° (from benzene). The same products but in reduced amt.s, were obtained when the condensation reaction was run in the presence of activated clay instead of

sulfamic acid. The structure of **II** was confirmed by an independent synthesis in which 4 g. *p*-*tolyl* was converted into the **K** deriv. and the latter was heated 1.5 hrs. to 230° with 1-diphenylmethylenec-2-chloro-3-phenylindene in the presence of 0.02 g. powd. Cu⁺ in addition to a product, m. 218°, identical with **II**; above, there was obtained some *1-diphenylmethylenec-3-phenylindene*, m. 204° (from Me₂CO). **III** (0.6 g.), 5 g. KOH, and 30 cc. 92% EtOH were refluxed 3 hrs. with stirring and the products were fractionally crystd. from ligroin, yielding a small amt. of an un-identified product, m. 215°, which did not have OH groups and which on heating above its m.p. was transformed to a compd. m. 177-80°, which was the principal product of the reaction and was identified as *1-diphenylmethylenec-2-(p-hydroxy-5-methylphenyl)-3-phenylindene*, m. 177-81° (from ligroin); treatment with Me₂SO₂ in the



presence of 30% NaOH gave the *Me ether*, m. 199° (from Me₂CO-petr. ether).

G. M. Kosolapoff

ZAL'KIND, YU. S.


PA 2/T4

USER/Chemistry - Hydrofluoric Acid Nov 1947
Chemistry - Catalysis

"Anhydrous Hydrogen Fluoride as a Catalyst," Yu. S.
Zal'kind, $\frac{1}{2}$ p

"Priroda" No 9

Article treats the use of hydrogen fluoride in the condensation of saturated hydrocarbon "isostuctures" with olefin according to the equation:



ID

2/T4

CA

10

Addition of hydrogen to acetylene derivatives. XXXVIII. Ethers of tetraphenylbutyndiol and their catalytic hydrogenation. Yu. S. Zait'kina and P. S. Rataev (Pedagog. Inst., Leningrad). *J. Gen. Chem. (U.S.S.R.)*, 17, 1358-67 (1947) (in Russian); cf. *C.A.*, 40, 3093. — $[Ph_4C(OH)C_2CC(OMe)Ph_4]_n$ (I) and the corresponding EIO compd. (II) were prep'd. according to Arbusov (*C.A.*, 18, 2327). The Pt compd. (III) was prep'd. by heating 10 g. $[Ph_4C(OH)C_2]_n$ (IV) with 0.6 g. iodine and 115 cc. $PrOH$ for 2 hrs. and cooling; III, m. 92-3°, sepd. on cooling. I with CrO_3 in $AcOH$ gave a diketone, $(Ph_4)_2C_2O_2$, a dark oil, which gives a semicarbazone, m. 151-2°, and on heating with O_2 and NH_3 , in the presence of alc. HCl gives a quinalidine, m. 172-3° (from $EIOH$); with excess CrO_3 in $AcOH$ the ketone gave methylidiphenylacetic acid, m. 99-100°. Similarly, II on oxidation gave a diketone, 1,1,4,4-tetraphenyl-1,4-diketone, 3,3-butanedione, a dark oil (semicarbazone, m. 163-9° (from $EIOH$); phenylhydrazine, m. 137-8°; quinalidine, acetic acid, m. 114°. IV (6 g.), 2.7 g. iodine, 40 cc. benzene, and 20 cc. abs. $EtOH$ boiled 5 hrs. gave 8.4 g.

$[Ph_4C(OH)C_2]_n$, m. 210° (from $EtOH$ -ligroin), which reacts with $EtMgBr$ and does not give C_2H_2 with KOH . To 5 g. I in 50 cc. $CHCl_3$ was added over 8 hrs. 4 g. Br in 35 cc. $CHCl_3$; after 7 hrs. heating to 45-50°, the mixt. was cooled to give 4.2 g. $[Ph_4(C_2O_2)CCBr]_n$, m. 190-7° (from ligroin). II similarly gave the EIO analog, m. 199-200°, and III gave the PtO analog, m. 195°. I (0 g.) in 70 cc. $CHCl_3$ treated with 4 atoms Cl and allowed to stand overnight gave 6.15 g. $[Ph_4(C_2O_2)CCCl]_n$, m. 194-5° (from Et_2O). IV (10 g.), 75 cc. benzene, and 1.12 g. $NaOAc$, treated in the cold with 23 g. Ac_2O , allowed to stand 0.8 hr., heated 2 hrs. to 45-50°, poured into water, gave no extn. with Et_2O 8.8 g. $[Ph_4(C_2O_4)C_2CC(OH)Ph_4]_n$, m. 133°. In the presence of pyridine with excess $AcCl$ there was obtained the di-Ac deriv., m. 164-70° (from ligroin). I, II, and III failed to add II in the presence of Pd or Pt . The mono-Ac deriv. added 2 H at a rate slower than that of the free glycol in the presence of 5 mg. Pd per 0.01 mol. of the substance at atm. pressure; the reaction stopped after addn. of 2 H. The di-Ac deriv. failed to add II either with Pd or Pt catalyst. The mono-Ac deriv. added 4 H with a Pt catalyst, giving after 19 hrs. (with 20 mg. Pt per 0.01 mole) the taad. compd., m. 157° (from $EtOH$); the product of addn. of 2 H m. 149°.

G. M. Korolapoff

ASA-SEA METALLURGICAL LITERATURE CLASSIFICATION

TECHNIQUE

140300-74

140300-1407 CRYO-OSC

140300-1407 CRYO-OSC

ZAL'KIND, Yu. S. DECEASED

12 to 15 May 1948, Moscow, first conference was held on history of Soviet chemistry, convened by Commission on the History of Chemistry, Acad. Sci. USSR. Many papers were presented by (ostensibly) members of this Commission.

"D. I. Mendeleev's 'Organic Chemistry'--First Russian Text on Organic Chemistry" (Issled 23 Nov 1948)

"Materials on the History of Soviet Chemical Science," published by Acad Sci USSR in Moscow-Leningrad 1950.

ZAL'KIND, YU. S.

"Synthesis of Long Chains of the Polypeptide Type," Priroda, No. 2, 1948.

ZAL'KIND, YU. S.

"Light Sensitive Glass," Priroda, No. 5, 1948. Prof.

ZAL'KIND, YU. S.

IA 9/49T43

USSR/Chemistry - Aldehydes
Chemistry - Synthesis

Sep 48

"New Method for Synthesizing Aldehydes," Yu. S.
Zal'kind, 1 p

"Priroda" No 9

Discussion of synthesis of acids from unsaturated compounds of hydrogen oxides and water was published in "Priroda" No 3, 1947. Further research reveals that if unsaturated derivatives and hydrogen oxides are treated with hydrogen in presence of a catalyst and heat, aldehydes can be produced (US Patent 2327066). Traces chemical process from formation of ethylene to final product--aldehyde.

9/49T43

ZAL'KIND, YU. S.

"Metallic Titanium and Zirconium," Priroda, No. 10, 1948.

ZAL'KIND, Yu. S. (Prof)

PA 25/49T87

USSR/Molecular Physics -- Radioactivity
Chemistry -- Periodic System Dec 48

"Review of A. E. Fersman's Book, 'The Role
of Mendeleev's Periodic Law in Contemporary
Science,'" Prof Yu. S. Zal'kind, 1 p

"Priroda" No 12

Brochure is series of lectures given by
Fersman during 1946, with well-organized and
thorough treatment of subject. Main considera-
tion is revolution which occurred in chemistry
with discovery of radioactive substances.
Recommended reading for the expert and for
the general public. Moscow, 1947, 56 pages.

25/49T87

price 2 rubles, 40 kopecks.

25/49T87

CH

XXXVIII. An erythritol of the diacetylene series and its catalytic hydrogenation. Yu. Zal'kina and L. Zhuravleva. *Zhur. Obshchey Khim.* [U. S. S. R. Chem.] 18, 984-9 (1948); cf. *C. A.* 42, 4101c. — Pt-Mafite (from 50 g. EtOH) and 10 g. $\text{Me}_2\text{C}(\text{OH})_2\text{CH}_2$ in EtOH were allowed to react with cooling and after standing overnight were treated with 12 g. benzil in EtOH, stirred several hrs., and allowed to stand 2 days, yielding cryst. $[\text{Pt}(\text{HO})_2\text{Me}_2\text{C}(\text{O})_2(\text{OH})_2]$, 60.0%, m. 144-5° (from EtOH), best sol. in Me_2CO , less sol. in EtOH or EtOAc ; heating with $\text{EtOH}-\text{Na}$ yields C_6H_6 ; addn. of Br in CHCl_3 was unsuccessful; oxidation by KMnO_4 gave $\text{Me}_2\text{C}(\text{OH})\text{CO}_2\text{H}$ and Br_2OH , as well as a ketone (semicarbazone, m. 181-2°). The product (3 g.) and 10 g. Ac_2O in the presence of a little HgPO_4 gave in 10 days at 20-5° 55.4% **c** *cryst. tetracetate*, m. 150-3° (from Me_2CO); the ale. with Ac_2O in the presence of NaOAc and pyridine gave the *diacetate*, m. 132-3°. Hydrogenation of the ale. in EtOAc over Pt is very slow after the addn. of sufficient H to convert the triple bonds to double bonds (essentially stopped after 30 min.), yielding *2,9-dimethyl-5,6-diphenoxy-3,7-deadiene-2,5,6,9-tetrol* (**D**), m. 146.5-47°. Similar hydrogenation over Pt black gives rapid and complete reaction, yielding the product as a noncrystallizable oil after addn. of 8 H atoms per mole. Hydrogenation of the tetracetate with the ale. itself, yielding the corresponding *tetracetate*, m. 148.5°, although after addn. of 4 H atoms the reaction still showed signs of slow progress. With a Pt catalyst the addn. of H is complete and proceeds further, apparently by replacement of the ester groupings by H, yielding hydrocarbons; this product was a noncrys. oil.

G. M. Kosolapoff

ASR-SLA METALLURGICAL LITERATURE

BIBLIOGRAPHY

SEARCHED MAY 1964

ECONOMIC

BULLETIN

PA 0747 21+

ZAL'KIND, YU. S.

USER/Chemistry - Organic Compounds
Chemistry - Halogenation

May 48

"The Action of Bromine and Chlorine on a Glycol of
the Diacetylene Series," Yu. S. Zal'kind and
N. N. Mel'teva, Lab of Org Chem, Leningrad Tech-
nological Inst imeni Lensoviet, 104 pp

"Zhur Obshch Khim" Vol XVIII (LXXX), No 5

Studies the action of bromine and chlorine on 2,
5-dimethyl-octa-3, 5-diene-2, 7-diol, which produces
a complex mixture of products. When they are
halogenated by a 10% alkali solution, 3, 4-dichlor
and 3, 4-dibrom-5-keto-2, 5-dihydrofuran is obtained.
Submitted 13 Sep 1947.

8/4971

CA

Action of bromine and chlorines on glycols of the acetates series. Yu. S. Zal'skikh and N. N. Mel'teva. Zhur. Obshch. Khim. 18, (905-1000)(1948)(in Russian). *3,5-Dimethyl-3,5-hexadiyne-2,7-diol* (**I**) with Cl or Br gives a complex mixt. of products. When 1 (m. 132.5%) of **I** and Atrikovich, t. 4.31, 428.0° (2.7 g.) in 50 ml. CHCl₃ was treated at -2° to -8° with 0.80 ml. Br₂, no HBr evolution was noted, but on attempts to remove the solvent by mild heating the product lost HBr and formed a black mass; removal of solvent in vacuo without heating gave viscous, yellow, liquid *di bromide*, which could not be distilled. A similar reaction, but using 4 atoms Br at 0-12°, led to partial loss of HBr in 2-3 hr., standing, and the removal of the solvent in *tetra- and hexabromides*. When 10 g. **I** in 50 cc. CHCl₃ was treated with 39.6 g. Br₂, the residue, after the removal of solvent, gave 1.2 g. solid, m. 167.9° (from EtOAc), corresponding to *C₁₀H₁₀OBr*; this substance is better obtained by the direct action of Br vapor on **I**; it is given the provisional formula: *C₁₀(CBr₂CHMe₂O)CBr₂CH₂CH(CBr₂CHMe₂O)Br*, since on oxidation by Cr oxide in AcOH it yields a

product, C₁₀H₁₀O₂, m. 129-0.5°, which is also obtained by the action of 10% NaOH on the crude bromide. Small amounts of Br-containing products, m. 101-37, 137.8°, and 70.8° were obtained but were not identified. The heptabromide is quite resistant to hydrolysis by 10% NaOH, as no reaction occurs in 24 hrs. at room temp., and only a slow reaction at 50°. Hydrolysis is 85% complete in 12 hrs. at 80-90°. Addition of 2.5 g. Cl to 5 g. **I** in CHCl₃ at 5-6° yields 2.1 g. unreacted **I** and an oil, b.p. 61-130°, which loses HCl slowly on distn., and appears to be a *tetra- and di- dichloride*; when 4 atoms Cl were used, 5.0 g. **I** gave 5 g. **I** and 2.0 g. oil, b.p. 108-109°, apparently a *di- and tetra- chloride*, m. 161. Use of 8 atoms Cl gave an undistillable yellow oil mixt. of *tetra- and hexa-Cl derivatives*. An excess of Cl passed at 0° through 100 cc. CHCl₃ gave a mixt. of products which decompl. on distn., and only 1 definite product, b. 62-7° in nitro (not specified), m. 122.3°, was obtained; this was C₁₀H₁₀OCl₆, identified as *dihexabromone*; washing the product with H₂O and 10% NaOH gave a slowly crystallizing oil, m. 81° (from ligroin), identified as *2,2-dimethyl-3,4-dichloro-5-keto-2,5-dihydrofuran*; the crude chlorination product oxidized with KMnO₄ gives dichloroacetone, HCOCl₂, and the above ketofuran. The formation of the furan derivs. is believed to take place through the cyclization of the *α,β-di-Cl* ketone intermediate. Possible reaction mechanisms are discussed. G. M. K.